

# Relative Positions in Finite Abelian Groups

Amritanshu Prasad

Institute Seminar Week 2011

# Spaces and Symmetry

# Spaces and Symmetry



Galileo



Galois



Jordan



Lie



Klein



Einstein

# Spaces and Symmetry



Galileo



Galois



Jordan



Lie



Klein



Einstein

Most physical (or mathematical) theories have two essential ingredients:

# Spaces and Symmetry



Galileo



Galois




Jordan



Lie



Klein



Einstein

Most physical (or mathematical) theories have two essential ingredients:

1. A space  $X$

# Spaces and Symmetry



Galileo



Galois



Jordan



Lie



Klein



Einstein

Most physical (or mathematical) theories have two essential ingredients:

1. A space  $X$
2. A symmetry group  $G$

# Spaces and Symmetry



Galileo



Galois



Jordan



Lie



Klein



Einstein

Most physical (or mathematical) theories have two essential ingredients:

1. A space  $X$
2. A symmetry group  $G$

The theory concerns quantities which are invariant under the symmetry group.

# Relative position

# Relative position

## Definition

Two pairs of points of  $X$  have the same relative position if  $G$  can transform one pair to the other:

# Relative position

## Definition

Two pairs of points of  $X$  have the same relative position if  $G$  can transform one pair to the other:

*$(x, y)$  and  $(x', y')$  have the same relative position if there exists  $g \in G$  such that  $x' = g \cdot x$  and  $y' = g \cdot y$ .*

# Relative position

## Definition

Two pairs of points of  $X$  have the same relative position if  $G$  can transform one pair to the other:

*$(x, y)$  and  $(x', y')$  have the same relative position if there exists  $g \in G$  such that  $x' = g \cdot x$  and  $y' = g \cdot y$ .*

## Simplest example

When  $X = \mathbf{R}$  and  $G = \mathbf{R}$  (acting by translations), the relative position of two points is determined by their difference:

# Relative position

## Definition

Two pairs of points of  $X$  have the same relative position if  $G$  can transform one pair to the other:

*$(x, y)$  and  $(x', y')$  have the same relative position if there exists  $g \in G$  such that  $x' = g \cdot x$  and  $y' = g \cdot y$ .*

## Simplest example

When  $X = \mathbf{R}$  and  $G = \mathbf{R}$  (acting by translations), the relative position of two points is determined by their difference:

*$(x, y)$  and  $(x', y')$  have the same relative position if and only if  $y - x = y' - x'$ .*

# Example: Euclidean Geometry

## Example: Euclidean Geometry

▶  $X = \mathbf{R}^n$

## Example: Euclidean Geometry

- ▶  $X = \mathbf{R}^n$
- ▶  $G = \mathbf{R}^n \rtimes SO(n)$

## Example: Euclidean Geometry

- ▶  $X = \mathbf{R}^n$
- ▶  $G = \mathbf{R}^n \rtimes SO(n)$

Relative position in Euclidean geometry

## Example: Euclidean Geometry

- ▶  $X = \mathbf{R}^n$
- ▶  $G = \mathbf{R}^n \rtimes SO(n)$

### Relative position in Euclidean geometry

The relative position of a pair of points in Euclidean geometry is determined by the distance between them:

## Example: Euclidean Geometry

- ▶  $X = \mathbf{R}^n$
- ▶  $G = \mathbf{R}^n \rtimes SO(n)$

### Relative position in Euclidean geometry

The relative position of a pair of points in Euclidean geometry is determined by the distance between them:

*The pairs  $(x, y)$  and  $(x', y')$  have the same relative position if and only if  $d(x, y) = d(x', y')$ .*

## Example: Euclidean Geometry

- ▶  $X = \mathbf{R}^n$
- ▶  $G = \mathbf{R}^n \rtimes SO(n)$

### Relative position in Euclidean geometry

The relative position of a pair of points in Euclidean geometry is determined by the distance between them:

*The pairs  $(x, y)$  and  $(x', y')$  have the same relative position if and only if  $d(x, y) = d(x', y')$ .*

Thus, the fundamental invariant of Euclidean geometry, namely distance, is a complete invariant of relative position.

# Points in a finite abelian group

# Points in a finite abelian group

- ▶  $X$  is a finite abelian group

# Points in a finite abelian group

- ▶  $X$  is a finite abelian group
- ▶  $G = X \rtimes \text{Aut}(X)$

# Points in a finite abelian group

- ▶  $X$  is a finite abelian group
- ▶  $G = X \rtimes \text{Aut}(X)$

In this case, the relative positions are combinatorial:

# Points in a finite abelian group

- ▶  $X$  is a finite abelian group
- ▶  $G = X \rtimes \text{Aut}(X)$

In this case, the relative positions are combinatorial:

[Dutta and Prasad, 2011](#)

We constructed a finite distributive lattice  $\Lambda$  and a function  $w : A \times A \rightarrow \Lambda$  such that  $(x, y)$  and  $(x', y')$  have the same relative position if and only if  $w(x, y) = w(x', y')$ .

# Points in a finite abelian group

- ▶  $X$  is a finite abelian group
- ▶  $G = X \rtimes \text{Aut}(X)$

In this case, the relative positions are combinatorial:

[Dutta and Prasad, 2011](#)

We constructed a finite distributive lattice  $\Lambda$  and a function  $w : A \times A \rightarrow \Lambda$  such that  $(x, y)$  and  $(x', y')$  have the same relative position if and only if  $w(x, y) = w(x', y')$ .

- ▶  $\Lambda$  depends on  $A$  only through the combinatorial invariants of its structure (and not the specific primes dividing its order)

# Subspaces in a vector space

# Subspaces in a vector space

- ▶  $X =$  the set of  $k$ -dimensional subspaces of  $\mathbf{R}^n$

# Subspaces in a vector space

- ▶  $X =$  the set of  $k$ -dimensional subspaces of  $\mathbf{R}^n$   
(Grassmannian space)

# Subspaces in a vector space

- ▶  $X =$  the set of  $k$ -dimensional subspaces of  $\mathbf{R}^n$   
(Grassmannian space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

# Subspaces in a vector space

- ▶  $X =$  the set of  $k$ -dimensional subspaces of  $\mathbf{R}^n$   
(Grassmannian space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

The relative position of a pair of subspaces is determined by the dimension of their intersection.


# Subspaces in a vector space

- ▶  $X =$  the set of  $k$ -dimensional subspaces of  $\mathbf{R}^n$   
(Grassmannian space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

The relative position of a pair of subspaces is determined by the dimension of their intersection.

*The pairs  $(V, W)$  and  $(V', W')$  have the same relative position if and only if  $\dim(V \cap V') = \dim(W \cap W')$ .*

This can be visualized as follows:




This can be visualized as follows:

$$V \cap W$$




$$V' \cap W'$$


This can be visualized as follows:



This can be visualized as follows:



This can be visualized as follows:



# Sugroups of a finite abelian group

# Sugroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

# Sugroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$

# Subgroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$
- ▶  $G = \text{Aut}(A)$

# Subgroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$
- ▶  $G = \text{Aut}(A)$

If pairs of subgroups  $(V, W)$  and  $(V', W')$  have the same relative position, then  $V \cap W$  and  $V' \cap W'$  are isomorphic.

# Sugroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$
- ▶  $G = \text{Aut}(A)$

If pairs of subgroups  $(V, W)$  and  $(V', W')$  have the same relative position, then  $V \cap W$  and  $V' \cap W'$  are isomorphic.

Unlike the case of subspaces of a vector space, the converse is not true.

# Subgroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$
- ▶  $G = \text{Aut}(A)$

If pairs of subgroups  $(V, W)$  and  $(V', W')$  have the same relative position, then  $V \cap W$  and  $V' \cap W'$  are isomorphic.

Unlike the case of subspaces of a vector space, the converse is not true.

Even if  $V \cap W$  is isomorphic to  $V' \cap W'$ , an isomorphism  $V \cap W \rightarrow V' \cap W'$  may not extend to an isomorphism  $V \rightarrow V'$  or  $W \rightarrow W'$ .

# Sugroups of a finite abelian group

Let  $A$  be a finite abelian group and  $B$  be a subgroup.

- ▶  $X =$  set of subgroups of  $A$  which are isomorphic to  $B$
- ▶  $G = \text{Aut}(A)$

If pairs of subgroups  $(V, W)$  and  $(V', W')$  have the same relative position, then  $V \cap W$  and  $V' \cap W'$  are isomorphic.

Unlike the case of subspaces of a vector space, the converse is not true.

Even if  $V \cap W$  is isomorphic to  $V' \cap W'$ , an isomorphism  $V \cap W \rightarrow V' \cap W'$  may not extend to an isomorphism  $V \rightarrow V'$  or  $W \rightarrow W'$ .

In general, this problem is known to be a *wild classification problem*.

# Nested families of subspaces in a vector space

## Nested families of subspaces in a vector space

▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$

## Nested families of subspaces in a vector space

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$  (Flag space)

## Nested families of subspaces in a vector space

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$  (Flag space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

# Nested families of subspaces in a vector space

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$  (Flag space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Suppose  $V = (V_0 \subset \cdots \subset V_n)$  and  $W = (W_0 \subset \cdots \subset W_n)$  is a pair of flags.

# Nested families of subspaces in a vector space

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$  (Flag space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Suppose  $V = (V_0 \subset \cdots \subset V_n)$  and  $W = (W_0 \subset \cdots \subset W_n)$  is a pair of flags.

In analogy with the Grassmannian space, we can compute dimensions of intersections:

$$d_{ij} = \dim(V_i \cap W_j)$$

## Nested families of subspaces in a vector space

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset \mathbf{R}^n \mid \dim(V_i) = i\}$  (Flag space)
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Suppose  $V = (V_0 \subset \cdots \subset V_n)$  and  $W = (W_0 \subset \cdots \subset W_n)$  is a pair of flags.

In analogy with the Grassmannian space, we can compute dimensions of intersections:

$$d_{ij} = \dim(V_i \cap W_j)$$

The relative position of two flags is determined by the matrix  $(d_{ij})$ .

The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:


Take  $n = 3$  and consider a pair of flags of subspaces:

$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$

The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:

$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$

$$V_1 \cap W_1$$

The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:


$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



The process of extending isomorphisms (as in the case of Grassmannian space) can be visualized as follows:

Take  $n = 3$  and consider a pair of flags of subspaces:

$$V_1 \subset V_2 \subset V_3 \subset V_4 \quad \text{and} \quad W_1 \subset W_2 \subset W_3 \subset W_4$$



# Organising the $d_{ij}$

## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

The dimension of successive subquotients is:

## Organising the $d_{ij}$


Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

The dimension of successive subquotients is:

$$w_{ij} := \dim \left( \frac{(V_i \cap W_j)/(V_i \cap W_{j-1})}{(V_{i-1} \cap W_j)/(V_{i-1} \cap W_{j-1})} \right) = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$



## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

The dimension of successive subquotients is:

$$w_{ij} := \dim \left( \frac{(V_i \cap W_j)/(V_i \cap W_{j-1})}{(V_{i-1} \cap W_j)/(V_{i-1} \cap W_{j-1})} \right) = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

The dimension of successive subquotients is:

$$w_{ij} := \dim \left( \frac{(V_i \cap W_j)/(V_i \cap W_{j-1})}{(V_{i-1} \cap W_j)/(V_{i-1} \cap W_{j-1})} \right) = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

Therefore, for each  $j$ , there exists unique  $i$  such that  $w_{ij} = 1$  (for all other values of  $i$ ,  $w_{ij} = 0$ ).

## Organising the $d_{ij}$

Each space  $W_j/W_{j-1}$  is one dimensional and

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces.

The dimension of successive subquotients is:

$$w_{ij} := \dim \left( \frac{(V_i \cap W_j)/(V_i \cap W_{j-1})}{(V_{i-1} \cap W_j)/(V_{i-1} \cap W_{j-1})} \right) = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

Therefore, for each  $j$ , there exists unique  $i$  such that  $w_{ij} = 1$  (for all other values of  $i$ ,  $w_{ij} = 0$ ).

Similarly, for each  $i$ , there is a unique  $j$  such that  $w_{ij} = 1$ .

Thus  $w = (w_{ij})$  is a permutation matrix.

# Bruhat Decomposition Theorem

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

This decomposition

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

This decomposition

- ▶ Plays a key role in the representation theory of general linear groups

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

This decomposition

- ▶ Plays a key role in the representation theory of general linear groups
- ▶ Algebraic geometry and cohomology theory of flag spaces

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

This decomposition

- ▶ Plays a key role in the representation theory of general linear groups
- ▶ Algebraic geometry and cohomology theory of flag spaces
- ▶ The theory of symmetric polynomials

# Bruhat Decomposition Theorem

## Relative positions of flags

The relative position of  $(V, W)$  is completely determined by  $w$ , where

$$w_{ij} = d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1},$$

which ranges over the set of  $n!$  permutation matrices.

This decomposition

- ▶ Plays a key role in the representation theory of general linear groups
- ▶ Algebraic geometry and cohomology theory of flag spaces
- ▶ The theory of symmetric polynomials
- ▶ Combinatorial identities involving Gaussian binomial coefficients

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

$$\blacktriangleright X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$$

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Given a pair of flags

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Given a pair of flags

$$V = (V_0 \subset \cdots \subset V_n) \quad \text{and} \quad W = (W_0 \subset \cdots \subset W_n),$$

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Given a pair of flags

$$V = (V_0 \subset \cdots \subset V_n) \quad \text{and} \quad W = (W_0 \subset \cdots \subset W_n),$$

we may try to measure their relative position by considering the isomorphism classes of the subgroups  $V_i \cap W_j$  of  $R^n$ .

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Given a pair of flags

$$V = (V_0 \subset \cdots \subset V_n) \quad \text{and} \quad W = (W_0 \subset \cdots \subset W_n),$$

we may try to measure their relative position by considering the isomorphism classes of the subgroups  $V_i \cap W_j$  of  $R^n$ .

This fails to determine relative position.

# Beyond the Bruhat decomposition

Let  $R = \mathbf{Z}/p^k\mathbf{Z}$ .

Consider

- ▶  $X = \{V_0 \subset V_1 \subset \cdots \subset V_n \subset R^n \mid V_i \cong R^i\}$
- ▶  $G = GL(n, \mathbf{R})$ , the group of invertible matrices

Given a pair of flags

$$V = (V_0 \subset \cdots \subset V_n) \quad \text{and} \quad W = (W_0 \subset \cdots \subset W_n),$$

we may try to measure their relative position by considering the isomorphism classes of the subgroups  $V_i \cap W_j$  of  $R^n$ .

This fails to determine relative position.

The strategy of systematically extending isomorphisms of subspaces fails because even if two finite abelian groups are isomorphic, not every isomorphism of subgroups extends to an isomorphism of the groups.

# Intersection matrices

## Intersection matrices

We still have

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces of  $W_i/W_{i-1} \cong R$ .

## Intersection matrices

We still have

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces of  $W_i/W_{i-1} \cong R$ .

But now,  $R = \mathbf{Z}/p^k\mathbf{Z}$  has subgroups, so several of the inclusions can be strict.

## Intersection matrices

We still have

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces of  $W_i/W_{i-1} \cong R$ .

But now,  $R = \mathbf{Z}/p^k\mathbf{Z}$  has subgroups, so several of the inclusions can be strict.

All we know is that if  $d_{ij} = \log_p |V_i \cap W_j|$ , then

$$w_{ij} := d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

## Intersection matrices

We still have

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces of  $W_i/W_{i-1} \cong R$ .

But now,  $R = \mathbf{Z}/p^k\mathbf{Z}$  has subgroups, so several of the inclusions can be strict.

All we know is that if  $d_{ij} = \log_p |V_i \cap W_j|$ , then

$$w_{ij} := d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

is a matrix with non-negative integer entries whose row and column sums are  $k$ .

## Intersection matrices

We still have

$$0 = \frac{V_0 \cap W_j}{V_0 \cap W_{j-1}} \subset \frac{V_1 \cap W_j}{V_1 \cap W_{j-1}} \subset \dots \subset \frac{V_n \cap W_j}{V_n \cap W_{j-1}} = \frac{W_j}{W_{j-1}}$$

is a nested sequence of subspaces of  $W_i/W_{i-1} \cong R$ .

But now,  $R = \mathbf{Z}/p^k\mathbf{Z}$  has subgroups, so several of the inclusions can be strict.

All we know is that if  $d_{ij} = \log_p |V_i \cap W_j|$ , then

$$w_{ij} := d_{ij} - d_{i-1,j} - d_{i,j-1} + d_{i-1,j-1}$$

is a matrix with non-negative integer entries whose row and column sums are  $k$ .

This is called the matrix of **intersection numbers** (Onn, Prasad and Vaserstein, 2006)

# Research problems

# Research problems

- ▶ Determine which integer matrices with row and column sums equal to  $k$  can occur as intersection numbers

# Research problems

- ▶ Determine which integer matrices with row and column sums equal to  $k$  can occur as intersection numbers
- ▶ How many relative positions can have the same intersection numbers (growth relative to  $p$ )

# Research problems

- ▶ Determine which integer matrices with row and column sums equal to  $k$  can occur as intersection numbers
- ▶ How many relative positions can have the same intersection numbers (growth relative to  $p$ )
- ▶ Study intersection numbers in the context of the rich combinatorial theory of integer matrices (Stanley's solution to ADG conjecture, RSK correspondence, etc.)

# Research problems

- ▶ Determine which integer matrices with row and column sums equal to  $k$  can occur as intersection numbers
- ▶ How many relative positions can have the same intersection numbers (growth relative to  $p$ )
- ▶ Study intersection numbers in the context of the rich combinatorial theory of integer matrices (Stanley's solution to ADG conjecture, RSK correspondence, etc.)
- ▶ Make the connections with the representation theory of  $GL_n(R)$