

Vectorlike Quarks at the LHC

Shrihari Gopalakrishna



Institute of Mathematical Sciences (IMSc), Chennai, India

with T.Mandal, S.Mitra, R.Tibrewala [arXiv:1107.4306, PRD84 (2011) 055001]

with T.Mandal, S.Mitra, G.Moreau [arXiv:1306.2656]

HIGGS-BSM 2013, ICTP Trieste, June 2013

Talk Outline

- Vector-like fermions we study
 - $b'_{(-1/3)}, t'_{(2/3)}, \chi_{(5/3)}$ ($t' \equiv T$)
 - Warped-space (Randall-Sundrum) model
 - Model independent whenever possible
- LHC Double and Single Resonant channels
 - Identify promising channels
 - Find luminosity required for discovery

Vectorlike ψ

- Vectorlike fermions:
 - both χ and χ^c present in the theory
 - can write vectorlike mass term $\mathcal{L} \supset M \chi\chi^c + h.c.$

Vectorlike fermions	Chiral (4-gen) fermions
M ok with Gauge Symmetry	M only after EWSB i.e. $\langle H \rangle$
can be arbitrarily heavy	Landau pole in Yukawa coupling
CC + NC tree-level decays	only CC tree-level decays
loops decoupling	some loops nondecoupling

EWSB induced mixing \implies Tree-level NC Couplings

Consider $t' \equiv T$ and b'

- $T_L b_L W^\pm$ and $b'_L t_L W^\pm$ CC couplings
- In Yukawa coupling $\langle \Sigma \rangle = v \implies t \leftrightarrow T, b \leftrightarrow b'$ mixing

$$\bullet \mathcal{L}_{\text{mass}} \supset \begin{pmatrix} t_L & t'_L \end{pmatrix} \begin{pmatrix} m_t & 0 \\ \tilde{m} & M_T \end{pmatrix} \begin{pmatrix} t_R \\ t'_R \end{pmatrix} + \\ \begin{pmatrix} b_L & b'_L \end{pmatrix} \begin{pmatrix} m_b & 0 \\ \tilde{m}_b & M_{b'} \end{pmatrix} \begin{pmatrix} b_R \\ b'_R \end{pmatrix} + h.c.$$

- leads to NC couplings $t'tZ$, $t'th$ and $b'bZ$, $b'bh$ also

EWSB induced mixing \implies Tree-level NC Couplings

Consider $t' \equiv T$ and b'

- $T_L b_L W^\pm$ and $b'_L t_L W^\pm$ CC couplings
- In Yukawa coupling $\langle \Sigma \rangle = v \implies t \leftrightarrow T, b \leftrightarrow b'$ mixing

$$\bullet \mathcal{L}_{\text{mass}} \supset \begin{pmatrix} t_L & t'_L \end{pmatrix} \begin{pmatrix} m_t & 0 \\ \tilde{m} & M_T \end{pmatrix} \begin{pmatrix} t_R \\ t'_R \end{pmatrix} + \\ \begin{pmatrix} b_L & b'_L \end{pmatrix} \begin{pmatrix} m_b & 0 \\ \tilde{m}_b & M_{b'} \end{pmatrix} \begin{pmatrix} b_R \\ b'_R \end{pmatrix} + h.c.$$

- leads to NC couplings $t'tZ, t'th$ and $b'bZ, b'bh$ also
- VL Tree-level Decays
 - $b' \rightarrow tW, b' \rightarrow bZ, b' \rightarrow bh$
 - $T \rightarrow bW, T \rightarrow tZ, T \rightarrow th$
 - $\chi \rightarrow tW$

Warped Model

SM in background 5D warped AdS space

[Randall, Sundrum '99]



$$ds^2 = e^{-2k|y|}(\eta_{\mu\nu} dx^\mu dx^\nu) + dy^2$$

Z_2 orbifold fixed points:

- Planck (UV) Brane
- TeV (IR) Brane

R : radius of Ex. Dim.

k : AdS curvature scale ($k \lesssim M_{pl}$)



Hierarchy prob soln:


- IR localized Higgs : $M_{EW} \sim ke^{-k\pi R}$: Choose $k\pi R \sim 34$
 - Gauge-theory dual is a composite Higgs model

Explaining SM mass hierarchy

Bulk Fermions explain SM mass hierarchy [Gherghetta, Pomarol 00][Grossman, Neubert '00]

$$\mathcal{L}_{Yuk}^{(5)} \supset \sqrt{|g|} \{ c_L k \bar{\psi}_L \psi_L + c_R k \bar{\psi}_R \psi_R + (\lambda_5 \bar{\psi}_R \psi_L H + h.c.) \}$$

$$\Psi_L(x, y) = \frac{e^{(2-c)ky}}{\sqrt{2\pi R N_0}} \Psi_L^{(0)}(x) + \dots$$



FCNC largely under control, but still strong constraints

AdS/CFT Correspondence

AdS/CFT Correspondence

[Maldacena, 1997]

- A classical supergravity theory in $AdS_5 \times S_5$ at weak coupling is **dual** to a 4D large-N CFT at strong coupling
- The CFT is at the boundary of AdS [Witten 1998; Gubser, Klebanov, Polyakov 1998]

$$Z_{CFT}[\phi_0] = e^{-\Gamma_{AdS}[\phi_0]}$$

$\mathcal{L} \supset \int d^4x \mathcal{O}_{CFT}(x) \phi_0(x)$ Eg: $\langle \mathcal{O}(x_1) \mathcal{O}(x_2) \rangle = \frac{\delta^2 Z_{CFT}[\phi_0]}{\delta \phi_0(x_1) \delta(x_2)}$ with Z_{CFT} given by the RHS	$\Gamma_{AdS}[\phi]$ supergravity eff. action $\phi(y, x)$ is a solution of the EOM ($\delta\Gamma = 0$) for given bndry value $\phi_0(x) = \phi(y = y_0, x)$
--	---

4D Duals of Warped Models

[Arkani-Hamed, Porrati, Randall, 2000; Rattazzi, Zaffaroni, 2001]

- Dual of Randall-Sundrum model **RS1 (SM on IR Brane)**
 - Planck brane \implies UV Cutoff; Dynamical gravity in the 4D CFT
 - TeV (IR) brane \implies IR Cutoff; Conformal invariance broken below a TeV
 - All SM fields are composites of the CFT
- Dual of Warped Models with **Bulk SM**
 - UV localized fields are elementary
 - IR localized fields (Higgs) are composite
 - 4D dual is Composite Higgs model [Georgi, Kaplan 1984]
 - Shares many features with Walking Extended Technicolor
 - Partial Compositeness
 - AdS dual is weakly coupled and hence calculable!
 - KK states are dual to composite resonances

Warped Bulk Gauge Group

[Agashe, Delgado, May, Sundrum '03]

Bulk gauge group : $SU(3)_{QCD} \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_X$

- Gauge Symmetry breaking:

- By Boundary Condition (BC):

$$A_{-+}(x, y) \text{ BC: } A|_{y=0} = 0; \partial_y A|_{y=\pi R} = 0$$

- $SU(2)_R \times U(1)_X \rightarrow U(1)_Y$

- By VEV of IR localized Higgs

$$\text{Higgs } \Sigma = (2, 2)_0$$

- $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$

Warped Fermions


- SM fermions : $(+, +)$ BC \rightarrow zero-mode
- “Exotic” fermions : $(-, +)$ BC \rightarrow No zero-mode
 - 1st KK vectorlike fermion

- Typical c_{t_R}, c_{t_L} : $(-, +)$ top-partners “light”

c : Fermion bulk mass parameter

[Choi, Kim, 2002] [Agashe, Delgado, May, Sundrum, 03]
 [Agashe, Perez, Soni, 04] [Agashe, Servant 04]

- Look for it at the LHC



[Dennis et al, '07] [Carena et al, '07] [Contino, Servant, '08]
 [Atre et al, '09, '11] [Aguilar-Saavedra, '09] [Mrazek, Wulzer, '09]
 [SG, Moreau, Singh, '10] [SG, Mandal, Mitra, Tibrewala, '11] [SG, Mandal, Mitra, Moreau : '13]

Fermion rep : $Zb\bar{b}$ not protected (DT model)

[Agashe, Delgado, May, Sundrum '03]

- Complete $SU(2)_R$ multiplet
 - $Q_L \equiv (\mathbf{2}, \mathbf{1})_{1/6} = (t_L, b_L)$
 - $\psi_{t_R} \equiv (\mathbf{1}, \mathbf{2})_{1/6} = (t_R, b')$
 - $\psi_{b_R} \equiv (\mathbf{1}, \mathbf{2})_{1/6} = (T, b_R)$
 - "Project-out" b' , T zero-modes by $(-, +)$ B.C.
 - New $\psi_{VL} : b', T$
- $b \leftrightarrow b'$ mixing
 - $Zb\bar{b}$ coupling shifted
 - So LEP constraint quite severe

Fermion rep : $Zb\bar{b}$ protected (ST & TT models)

- $Q_L = (2, 2)_{2/3} = \begin{pmatrix} t_L & \chi \\ b_L & T \end{pmatrix}$ [Agashe, Contino, DaRold, Pomarol '06]
- $Zb_L\bar{b}_L$ protected by custodial $SU(2)_{L+R} \otimes P_{LR}$ invariance
 $Wt_L b_L, Zt_L t_L$ not protected, so shifts

Fermion rep : $Zb\bar{b}$ protected (ST & TT models)

- $Q_L = (2, 2)_{2/3} = \begin{pmatrix} t_L & \chi \\ b_L & T \end{pmatrix}$ [Agashe, Contino, DaRold, Pomarol '06]
 - $Zb_L\bar{b}_L$ protected by custodial $SU(2)_{L+R} \otimes P_{LR}$ invariance
 $W_{t_L b_L}, Z_{t_L t_L}$ not protected, so shifts

Two t_R possibilities:

① Singlet t_R (ST Model) : $(1, 1)_{2/3} = t_R$ New $\psi_{VL} : \chi, T$

② Triplet t_R (TT Model) :

$$(1, 3)_{2/3} \oplus (3, 1)_{2/3} = \psi'_{t_R} \oplus \psi''_{t_R} = \begin{pmatrix} \frac{t_R}{\sqrt{2}} & \chi' \\ -\frac{t_R}{\sqrt{2}} & b' \end{pmatrix} \oplus \begin{pmatrix} \frac{t''}{\sqrt{2}} & \chi'' \\ b'' & -\frac{t''}{\sqrt{2}} \end{pmatrix}$$

New $\psi_{VL} : \chi, T, \chi', b', \chi'', t'', b''$

Yukawa Couplings

Yukawa Couplings

- No $Zb\bar{b}$ protection






$$\mathcal{L}_{\text{Yuk}} \supset \lambda_t \bar{Q}_L \Sigma \psi_{tR} + \lambda_b \bar{Q}_L \Sigma \psi_{bR} + h.c.$$

- With $Zb\bar{b}$ protection





- ST Model $\mathcal{L}_{\text{Yuk}} \supset \lambda_t \text{Tr}[\bar{Q}_L \Sigma] t_R + h.c.$

- TT Model $\mathcal{L}_{\text{Yuk}} \supset \lambda_t \text{Tr}[\bar{Q}_L \Sigma \psi'_{tR}] + \lambda'_t \text{Tr}[\bar{Q}_L \Sigma \psi''_{tR}] + h.c.$






Warped model b' parameters



Warped model χ parameters



Warped model t' parameters







b' Phenomenology at the LHC


[SG, T.Mandal, S.Mitra, R.Tibrewala, arXiv:1107.4306]

[SG, T.Mandal, S.Mitra, G.Moreau : arXiv:1306.2656]

Warped model b' : Γ and BR



b' Single & Double Resonant channels





... followed by $b_2 \rightarrow bZ$

- Both b_2 on-shell : **Double Resonant (DR)** channel
- Only one b_2 on-shell : **Single Resonant (SR)** channel
 - $|M(bZ) - M_{b_2}| \geq \alpha_{cut} M_{b_2}; \quad \alpha_{cut} = 0.05$

b' Double Resonant

Pair Production : $pp \rightarrow b'\bar{b}' \rightarrow bZ\bar{b}Z \rightarrow bjj\bar{b}ll$




Cuts:

- Rapidity:* $-2.5 < y_{b,j,Z} < 2.5$,
- Transverse momentum:* $p_{T,b,j,Z} > 25$ GeV,
- Invariant mass cuts:*
- $M_Z - 10$ GeV $< M_{jj} < M_Z + 10$ GeV,
- $0.95M_{b_2} < M_{(bZ)} < 1.05M_{b_2}$.

b' Single Resonant - I

Single Resonant : $bg \rightarrow b'bZ \rightarrow bZbZ \rightarrow bbJJ\ell\ell$

Model Independent LHC-14 reach





Brown dots : DT Model

Green dots : TT Model

b' Single Production - II

Single Production : $bg \rightarrow b'Z \rightarrow bZZ \rightarrow bj\ell\ell$




Cuts:


Rapidity: $-2.5 < y_{b,j,Z} < 2.5$,
 Transverse momentum: $p_{T,b,j,Z} > 0.1M_{b_2}$,
 Invariant mass cuts:
 $M_Z - 10 \text{ GeV} < M_{jj} < M_Z + 10 \text{ GeV}$,
 $0.95M_{b_2} < M_{(bZ)} \text{ OR } (bjj) < 1.05M_{b_2}$.

χ Phenomenology at the LHC

[SG, T.Mandal, S.Mitra, G.Moreau : arXiv:1306.2656]


Warped model Γ_χ 

ST Model



TT Model


χ Double and Single Resonant channels



$$pp \rightarrow \chi t W \rightarrow t W t W \rightarrow t W t \nu$$

X	M_χ (GeV)	σ_{tot} (fb)	σ_{SR} (fb)	cuts	S (fb)	BG (fb)	\mathcal{L} (fb^{-1})
X_1	500	2406	261.5	Basic	977.5	3.257	-
				Disc.	146.1	0.115	0.826
X_2	750	235.5	29.31	Basic	99.99	3.257	-
				Disc.	42.74	0.115	2.824
X_3	1000	39.19	5.198	Basic	17.92	3.257	-
				Disc.	11.36	0.115	10.63
X_4	1250	8.576	1.231	Basic	4.305	3.257	-
				Disc.	3.226	0.115	37.42
X_5	1500	2.188	0.364	Basic	1.235	3.257	-
				Disc.	1.010	0.115	119.5
X_6	1750	0.613	0.121	Basic	0.393	3.257	-
				Disc.	0.339	0.115	355.8

χ Single Resonant Channel






Blue Dots - ST Model Green Dots - TT Model

t' Phenomenology at the LHC


[SG, Tanumoy Mandal, Subhadip Mitra, Gregory Moreau : arXiv:1306.2656]

See also: [Harigaya et al., '12] [Giridhar, Mukhopadhyaya, 2012] [Azatov et al., '12]
[Berger, Hubisz, Perelstein, '12] [Cacciapaglia et al., '10, '12]

Warped model t' BR




t' Double and Single Resonant channels



$pp \rightarrow t_2 th \rightarrow thth \rightarrow tbbtbb \rightarrow 6 b 4 j$ (4 b-tags)

T	M_{t_2} (GeV)	σ_{tot} (fb)	σ_{SR} (fb)	cuts	S (fb)	BG (fb)	\mathcal{L} (fb^{-1})
T_1	500	1207	223.0	Basic	237.4	102.7	-
				Disc.	52.38	0.389	6.379
T_2	750	115.2	18.30	Basic	22.67	102.7	-
				Disc.	13.25	0.389	25.22
T_3	1000	18.38	2.715	Basic	3.088	102.7	-
				Disc.	2.421	0.389	138.0
T_4	1250	3.821	0.590	Basic	0.477	102.7	-
				Disc.	0.415	0.389	1889.2

t' Single Resonant channel



Blue Dots - ST Model Green Dots - TT Model

Conclusions

- Vector-like quarks
 - have NC (and CC) tree-level decays
 - could be much lighter than V_{μ}^{KK} in warped models
- Identified promising DR and SR channels
 - SR can probe EW couplings
- 14 TeV LHC with $\approx 300 \text{ fb}^{-1}$ reach about 1.5 - 2 TeV in DR
- New ATLAS & CMS results
 - limits around $M_{\psi} \gtrsim 750 \text{ GeV}$

BACKUP SLIDES

BACKUP SLIDES


b' Pair Production Details

$$pp \rightarrow b' \bar{b}' \rightarrow bZ\bar{b}Z \rightarrow bj\bar{j}b\ell\ell$$

M_{b_2} (GeV)	Signal σ_s (in fb)		Background σ_b (in fb)				\mathcal{L} (fb^{-1})
	$bZbZ$		$bZbZ$		$(bj\bar{j}bZ)_{\text{tot}}$		
	y, p_T cuts	All cuts	y, p_T cuts	All cuts	y, p_T cuts	All cuts	
250	25253	25082	21.804	0.3797	16938	29.52	0.021
500	171.34	148.69	21.804	0.047	16938	3.74	3.514
750	14.508	12.221	21.804	0.0097	16938	0.997	42.752
1000	2.314	1.9214	21.804	0.0027	16938	0.259	271.92
1250	0.484	0.399	21.804	0.0011	16938	0.048	1310

M_{b_2} (GeV)	QCD background (in fb)					
	$bjjbZ$		$bbjbZ$		$bbbbZ$	
	y, p_T cuts	All cuts	y, p_T cuts	All cuts	y, p_T cuts	All cuts
250	16790	27.304	255.41	2.7	81.01	1.92
500	16790	3.513	255.41	0.256	81.01	0.194
750	16790	0.958	255.41	0.031	81.01	0.057
1000	16790	0.2514	255.41	0.0052	81.01	0.008

b' Signature (Model Independent)



Benchmark Points (Model I):

M_{b_2} (GeV)	250	500	750	1000	1250	1500
$\kappa_{b_2 b Z}^L$	0.185	0.121	0.084	0.064	0.051	0.043
$\kappa_{b_2 t W}$	0.322	0.161	0.107	0.080	0.064	0.054
$\kappa_{hb_L b_2 R}$	0.714	0.937	0.972	0.985	0.990	0.993
M_{b_2} (GeV)	1750	2000	2250	2500	2750	3000
$\kappa_{b_2 b Z}^L$	0.037	0.032	0.029	0.026	0.024	0.022
$\kappa_{b_2 t W}$	0.046	0.040	0.036	0.032	0.029	0.027
$\kappa_{hb_L b_2 R}$	0.995	0.996	0.997	0.998	0.998	0.998

b' Single Resonant II Details

$$pp \rightarrow b'Z \rightarrow bZZ \rightarrow bjj\ell^+\ell^-$$

$M_{b'}$ (GeV)	signal σ_s (in fb)		background σ_b (in fb)				$\mathcal{L}^{\text{SemiLep}}$ (fb^{-1})
	$bjjZ$		$(bjjZ)_{EW}$		$(bjjZ)_{QCD}$		
	Primary cuts	all cuts	Primary cuts	all cuts	Primary cuts	all cuts	
250	1017.66	995.86	77.03	10.33	7853.02	867.82	0.66
500	16.84	15.50	8.81	0.68	419.75	14.11	45.94
750	1.26	1.14	1.85	0.10	56.26	0.86	551.26
1000	0.14	0.12	0.47	0.01	12.38	0.05	3399.67

$M_{b'}$ (GeV)	QCD background (in fb)		
	$bjjZ$	$bjbZ$	$bbbZ$
250	546.36	634.32	17.19
500	10.14	7.76	0.35
750	0.52	0.66	0.03
1000	0.02	0.06	0.002