

# Systems Biology: A Personal View

## XXV. Waves in Biology: Excitable Media

Sitabhra Sinha  
IMSc Chennai

Turing's model for pattern formation is a specific example of

# Reaction-Diffusion Equations

Activator:  $\partial u / \partial t = D_u \Delta u + f(u, v)$

Inhibitor:  $\partial v / \partial t = D_v \Delta v + g(u, v)$

For Turing patterns to occur,  $D_u < D_v$

But, what if  $D_u > 0$  and  $D_v = 0$



Reaction-Diffusion Media

Excitable  
Media

*Spatiotemporal  
Pattern formation in  
excitable media  
models*

# What is Excitable Media ?

**... that biology should be mindful of it ?**

- **pancreatic beta cells**
- **neurons**
- **cardiac myocytes**
- **pregnant uterus**

# What is Excitable Media ?

Think of an excitable person

**Resting** State


**Excited** State

Transition from resting to excited state if **stimulation** exceeds a **threshold**

Immediately after one excitation, the medium cannot be excited even by a very high stimulus for a **resting (refractory) period**


# A simple model

$$\frac{dx}{dt} = f(x) = x(x - \alpha)(1 - x) - y$$



$$\frac{dy}{dt} = \varepsilon (k x - y)$$

# Phase plane dynamics of the simple model



Below threshold  $\rightarrow$  decays to resting state

Above threshold  $\rightarrow$  excitation (large excursion from stable resting state).

This simple model is none other than the


# Fitzhugh-Nagumo model

$$\frac{du_e}{dt} = F_e(u_e, v) = Au_e(u_e - \alpha)(1 - u_e) - v,$$
$$\frac{dv}{dt} = \epsilon(u_e - v),$$

Developed by R Fitzhugh (1961) [who called it the Bonhoeffer-Van der Pol model] and J Nagumo (1962) to isolate the essential concepts of excitation propagation

**Richard Fitzhugh:** Simplified the Hodgkin-Huxley equations describing spike generation in squid giant axon

**Jin-ichi Nagumo:** built monostable multivibrator electronic circuit using tunnel (Esaki) diodes  
Esaki diodes have cubic I-V curve similar to that used in Fitzhugh's eqn



Tokyo U

# Spatial propagation of excitation waves

## Example: Cardiac tissue

functions as a syncytium: constituent cells are electrically synchronized through specialized channels called gap junctions

Delay in spread of depolarization to neighboring cells via gap junctions ... but usually the discrete nature of cardiac myocytes can be ignored and propagation of excitation assumed to be continuous.

Mathematically approximated as a **diffusion equation** yielding the partial differential equation

$$\frac{\partial V}{\partial t} = D \frac{\partial^2 V}{\partial x^2} - \frac{I_{ion}}{C_m}$$

with diffusion constant  $D = \frac{G_i}{S_v C_m}$

$G_i$ : Bulk intracellular conductivity  
 $S_v$ : Surface to volume ratio of cells  
 $C_m$ : Membrane capacitance





Biological tissue is anisotropic  $\Rightarrow D$  is a tensor

E.g., action potentials travel along the direction of muscle fibers much faster than transverse to it



# Microscopic anatomy of cardiac muscle


<http://antranik.org/>



(b)

# Gap junctions formed by assembly of connexin proteins

connexon: assembly of six connexin proteins forming the pore for a gap junction between cytoplasm of a pair of connected cells – allowing bidirectional flow of ions and signaling molecules



# Example: one-dimensional chain of cells

Nearest neighbors connected by gap junctions

$V_n$ : transmembrane potential of  $n$ -th cell

$I_n$ : current from the  $n$ -th to the  $(n-1)$ -th cell



The net current that passes through gap junctions of the  $n$ -th cell:

$$I_{junction} = I_n - I_{n+1} = g_{gap}(V_n - V_{n-1}) - g_{gap}(V_{n+1} - V_n)$$

where  $g_{gap}$ : gap-junction conductance

Using continuum approximation,


$$I_{junction} = -g_{gap} \frac{\partial^2 V}{\partial x^2}$$

yielding the partial differential equation describing spatial diffusion of excitation

$$C_m \frac{\partial V}{\partial t} = -I_{ion} - I_{junction} = g_{gap} \frac{\partial^2 V}{\partial x^2} - I_{ion}$$

# Propagating Waves in Excitable Media

- Excited cells can excite their neighboring cells via diffusion



- The propagating excitation waves can collide and annihilate each other

... resulting in spontaneous pattern formation, such as single or multiple spiral waves

