

Planar depth and planar subalgebras

Zeph Landau

Mathematical Sciences Research Institute

E-mail: landau@math.berkeley.edu

and

V.S. Sunder

The Institute of Mathematical Sciences, Chennai

E-mail: sunder@imsc.ernet.in

We consider the notion of *planar depth* of a planar algebra, *viz.*, the smallest n for which the planar algebra is generated by its ' n -boxes'. We establish a simple result which yields a sufficient condition, in terms of the principal graph of the planar algebra, for the planar depth to be bounded by k . This suffices to determine the planar depth of the E_6 , E_8 and the $\frac{5+\sqrt{13}}{2}$ subfactors.

We then consider a planar subalgebra of the 'group planar algebra' which is naturally associated with a group Θ of automorphisms of the given group G . We show that this planar algebra corresponds to the 'subgroup-subfactor' associated with the inclusion $\Theta \subset (G \rtimes \Theta)$ (given by the semi-direct product extension). We conclude with a discussion of the planar depth of this planar algebra P^Θ in some examples.

Key Words: operator algebras, subfactor, planar algebra, standard invariant.

1. INTRODUCTION

Jones defined the notion of a *planar algebra* in [6], where he showed, using Popa's characterisation in [10] of the so-called λ -lattices, that (spherical, connected C^* -) planar algebras are in bijection with standard invariants of extremal subfactors. In particular, each planar algebra is the planar algebra associated with at least one subfactor.

Given a planar algebra $P = \{P_n\}$, let us define P^n to be the planar subalgebra (of P) generated by P_n ; then we have a tower

$$P^0 \subseteq P^1 \subseteq P^2 \subseteq \dots \subseteq P^n \subseteq P^{n+1} \subseteq \dots \cup_{n=0}^{\infty} P^n = P \quad (1.1)$$

of planar subalgebras of P . (Note that P^0 is nothing but the Temperley-Lieb planar algebra TL - with the same δ as the given planar algebra.) It is fairly easy to see that this tower will stabilise at a finite stage *provided* the initial planar algebra P 'has finite depth'. Of course, this finite depth condition is far from necessary for the tower to stabilise; for instance, the Temperley-Lieb planar algebra TL - cf. [6] - satisfies $TL = (TL)^k, \forall k \geq 0$ (since it is generated by its zero boxes) although TL has infinite depth in the 'generic case'. We shall call the smallest integer for which the tower (1.1) stabilises the *planar depth*; i.e. P has planar depth $k, 0 \leq k \leq \infty$ if $P = P^l \Leftrightarrow l \geq k$. Since 'planar depth \leq depth', it is natural to ask what the planar depth of a finite-depth planar algebra can be.

In section 2, we prove a fairly simple fact (cf. Corollary 2.2.2 and Proposition 2.2.1) that suffices to determine the planar depth of some planar algebras of finite depth.

The first author (in [9]) gave a presentation of the planar algebra $P(G)$ of the group subfactor corresponding to the fixed-points of an outer action of a finite group G on a II_1 factor. He then considered the scenario of a finite group Θ acting on the group G as group-automorphisms. He showed (Theorem 9 of [9]) that the planar algebra that is generated by the elements in $P(G)_2 \equiv \mathbb{C}G$ which are fixed by (the linear extension of) Θ , is strictly smaller than $P(G)$. He discussed this subplanar algebra for several examples.

In section 3, we begin by observing that if Θ acts on G as above, then there is a natural associated action of Θ on the planar algebra $P(G)$. The invariants of this action, call it P^Θ (the group G and the action of Θ on it will be fixed once and for all), yield a planar subalgebra of $P(G)$. We see that the 'subplanar algebra' of the last sentence of the previous paragraph is just what we call $(P^\Theta)^2$.

We show that P^Θ can in fact be identified with the planar algebra associated with the subgroup-subfactor corresponding to the subgroup Θ of the semi-direct product $G \rtimes \Theta$ and in particular *has finite depth*. This finite depth statement implies, as observed earlier, that P^Θ has finite planar depth. We conclude by discussing some examples that illustrate several possible features of the tower (1.1).

2. A CRITERION FOR ESTIMATING PLANAR DEPTH

We shall use the terminology of [6] for planar algebras. Thus a planar algebra P is a tower $\{P_n : n = 0, 1, 2, \dots\}$ of finite-dimensional C^* -algebras equipped with 'an action of the coloured operad \mathcal{P} of labelled tangles'. We shall, as in [6], use the term ' k -boxes' to denote elements of P_k .

As stated in the introduction, if $P = \{P_n\}$ is a planar algebra, we shall denote by P^n the planar subalgebra of P generated by P_n . We shall denote by P_k^n the vector space of k -boxes in P^n .

PROPOSITION 2.2.1. *Suppose the principal graph Γ of P satisfies the following condition for some $n \in \mathbb{N}$:*

(†) *each vertex at distance $(n + 1)$ from $*$ is adjacent to a unique vertex at distance n from $*$, and these latter vertices (= neighbours) are all distinct.*

Then $P^n = P^{n+1}$.

Proof: The proof relies on an inspection of the tower

$$P_n^n \subset P_{n+1}^n \subseteq P_{n+1}^{n+1}, \quad (2)$$

showing that the condition (†) implies that the second inclusion is actually an equality, and thus $P^n = P^{n+1}$.

We introduce some notation for the Bratteli diagrams for the inclusions in (2):

(i) Denote by X , Y , and Z the vertex set for P_n^n , P_{n+1}^n , and P_{n+1}^{n+1} respectively.

(ii) Define the set $B =$

$$\{(x, y) : x \in X \text{ is connected to } y \in Y \text{ in the Bratteli diagram.}\} \\ \amalg \\ \{(y, z) : y \in Y \text{ is connected to } z \in Z \text{ in the Bratteli diagram.}\},$$

where \amalg denotes the disjoint union.

(iii) For sets S and T we shall write $(S, T) = Id$ if there is a bijection $f : S \rightarrow T$ such that for all $s \in S$, $t \in T$

$$(s, t) \in B \Leftrightarrow t = f(s).$$

(iv) Since the inclusion $P_n^n = P_n^n \subset P_{n+1}^n = P_{n+1}^n$ is part of a Jones tower, there is a natural partition of X and Z into ‘old’ and ‘new’ vertices (see [4] for this terminology) which we write as $X = X^O \amalg X^N$ and $Z = Z^O \amalg Z^N$.

(v) The inclusion $P_n^n \subset P_{n+1}^n$ is also part of a Jones tower and thus X and Y have a natural partition into ‘old’ and ‘new’ vertices as well. Furthermore, this partition for X is the same as the partition in (iv) above since the two Jones towers are identical up to $P_n^n = P_n^n$. We denote the partition of Y by $Y = Y^O \amalg Y^N$.

With this notation, showing equality between P_{n+1}^n and P_{n+1}^{n+1} is equivalent to showing $(Y, Z) = Id$. We shall accomplish this by showing the four statements:

$$\begin{aligned} (Y^N, Z^N) &= Id, & (Y^O, Z^O) &= Id, \\ (Y^N \times Z^O) \cap B &= \emptyset, & (Y^O \times Z^N) \cap B &= \emptyset. \end{aligned} \quad (3)$$

We have the following facts about the Bratteli diagram:

- (a) $(\{y\} \times Z) \cap B \neq \emptyset$ for all $y \in Y$.
- (b) The nature of the partition of X, Y and Z into old and new vertices implies that

- (i) $(X^O \times Y^N) \cap B = \emptyset$,
- (ii) for all $y \in Y^O$, $(X^O \times \{y\}) \cap B \neq \emptyset$,
- (iii) there is no $y \in Y$ such that

$$(X^O \times \{y\}) \cap B \neq \emptyset \text{ and } (\{y\} \times Z^N) \cap B \neq \emptyset.$$

- (c) It follows from (bii) and (biii) that $(Y^O \times Z^N) \cap B = \emptyset$.

(d) Since both Y^O and Z^O correspond to the set of minimal central projections in the ‘basic construction ideal’ $P_n e_n P_n$ (with $e_n \in P_{n+1}$ denoting the n -th Jones projection), we have $(Y^O, Z^O) = Id$ and $(Y^N \times Z^O) \cap B = \emptyset$.

Thus we have shown the last three statements in (3); it remains to show that $(Y^N, Z^N) = Id$. But given $(Y^N \times Z^O) \cap B = \emptyset$ and (a), this is just what condition (†) ensures. \square

COROLLARY 2.2.2. *Let P be a planar algebra with associated principal graph Γ . Suppose Γ has no double bonds, no vertices of degree greater than 3, and a unique vertex of degree 3. If this degree 3 vertex is of distance $(k - 1)$ from $*$, then P has planar depth k and is generated (as a planar algebra) by one k -box.*

Proof: It follows directly from Proposition 2.2.1 that $P^l = P^{l+1}$ for all $l \geq k$; hence the planar depth of P is finite and at most k . On the other hand, a simple dimension argument shows that $P^{k-1} = P^0$ is nothing but the Temperley-Lieb planar algebra TL and in particular $P^{k-1} \neq P^k$, so the planar depth is exactly k . Finally, the hypothesis shows that $\dim P_k = \dim TL_k + 1$, and this shows that any element of P_k which is not in P_k^{k-1} will generate P as a planar algebra. \square

REMARK 2.2.3. *Three remarks are in order here.*

(a) *Special cases to which this Corollary applies are the cases when Γ is D_{2n}, E_6, E_8 and the principal graph of the $\frac{5+\sqrt{13}}{2}$ subfactor of [1].*

(b) *Of the pair of principal graphs associated to the $\frac{5+\sqrt{13}}{2}$ subfactor of [1], one graph has a unique triple point, while the other has two triple points at different distances from $*$. (This feature also holds in each case of the heirarchy of pairs of graphs listed (see [5], case (2)) by Haagerup as other possible finite principal graphs of subfactors of index less than $3 + \sqrt{3}$.)*


Since the planar depth of a subfactor is the same as that of its dual, we see (as illustrated by the graph (in [5]), referred to above, with two triple points) that it is possible for a principal graph to have a triple point at a distance $(k - 1)$ from $$ and still have planar depth strictly smaller than k .*

(c) *As remarked above (and as can be seen from Jones' description of $P^{M \subset M_1}$), a planar algebra and its dual planar algebra have the same planar depth. This is not the case for the usual depth of a subfactor; the $\frac{5+\sqrt{13}}{2}$ subfactor provides an example where the depths of the subfactor and its dual differ. This may be cited as one reason why 'planar depth' is a more natural notion than the usual depth.*

3. SOME PLANAR SUBALGEBRAS OF THE GROUP PLANAR ALGEBRA

We shall only be concerned with planar algebras P which come equipped with a 'presentation (whose symbol Φ we shall suppress) by a collection $L = \coprod_{n=0}^{\infty} L_n$ of labels'. (Again see [6] for notation.)

In fact, following [9], we shall be primarily concerned with the planar algebra $P(G)$, which has a presentation as above with generators given by $L_2 = G$ and $L_k = \emptyset$ for $k \neq 2$, the relation that a simple closed loop (of either orientation) be the scalar $|G|^{\frac{1}{2}}$ and the additional six relations labelled 00,0,1,2,3,4 below (and in Theorem 5 of [9]). We shall assume that we are given a group Θ and an action $\alpha : \Theta \rightarrow \text{Aut}(G)$. Nothing is changed if we replace Θ by $\Theta/\ker \alpha$, so we assume that Θ acts faithfully, i.e., that α is 1-1.




We consider the map on tangles that replaces the label of each 2-box with the label's image under $\theta \in \Theta$. Since θ is a group automorphism, it is seen that the set of relations defining $P(G)$ is unchanged by this map and thus this map defines an automorphism of $P(G)$ which we shall continue to denote by θ . It follows that the set P^Θ of invariants for this action of Θ on $P(G)$ is a sub-planar algebra of P , and that the set of Θ -invariant k -boxes of $P(G)$ constitutes precisely the set of k -boxes of P^Θ .


For each $k = 1, 2, \dots$ and $\theta \in \Theta$, let $\alpha_\theta^{(k)} \in \text{Aut}(G^k)$ be defined by $\alpha_\theta^{(k)}(g_1, g_2, \dots, g_k) = (\alpha_\theta(g_1), \alpha_\theta(g_2), \dots, \alpha_\theta(g_k))$. When the context is clear, we shall simply write $\theta(g_1, \dots, g_k)$ for what we have defined above as $\alpha_\theta^{(k)}(g_1, \dots, g_k)$.

For convenience of reference, we shall gather various simple facts about bases for the spaces $P(G)_k$ in the form of the following remark.

REMARK 3.3.1. (a) It was shown in Theorem 6 of [9] that if we let $T(g_1, \dots, g_{k-1})$ denote the labelled k -tangle given by




for k odd, and




for k even, then $\{T(\bar{g}) : \bar{g} \in G^{k-1}\}$ is an orthonormal basis of $P(G)_k$ (with respect to the inner product given by the natural trace).

(b) We shall find it convenient to use a slightly different basis (which, as we shall see, is actually just a rearrangement of the basis in (a)): define $S(\bar{g}), \bar{g} \in G^{k-1}$ to be the labelled k -tangle given by



for k odd, and



for k even.

(c) On the one hand, the map $G^{k-1} \ni \bar{g} \mapsto \bar{h} \in G^{k-1}$ defined by

$$h_i = \begin{cases} g_i^{-1} g_{i+1} & \text{if } i < k-1 \\ g_{k-1}^{-1} & \text{if } i = k-1 \end{cases} \quad (4)$$

is clearly a bijection with inverse given by $\bar{h} \mapsto \bar{g}$, where $g_i = (h_i h_{i+1} \dots h_{k-1})^{-1}$.

On the other hand, it is an easy exercise to use relation 3. above to show that - with \bar{g} and \bar{h} related as above - we have $S(\bar{g}) = T(\bar{h})$, and hence that $\{S(\bar{g}) : \bar{g} \in G^{k-1}\}$ is an orthonormal basis of $P(G)_k$ as well.

(d) It follows from the definitions that $\{S(1, g_2, g_3, \dots, g_{k-1}) : (g_2, g_3, \dots, g_{k-1}) \in G^{k-2}\}$ is an orthonormal basis for $P(G)_{1,k}$. (*Reason:* Clearly $S(\bar{g}) \in P(G)_{1,k}$ if $g_1 = 1$, while relation 2. above shows that $S(\bar{g}) \perp P(G)_{1,k}$ if $g_1 \neq 1$.)

(e) We will also need to decompose an $S(\bar{g}), \bar{g} \in G^{k-2}$, when regarded as an element of $P(G)_k$ under the natural inclusion of $P(G)_{k-1}$ into $P(G)_k$, in terms of the basis $\{S(\bar{h}) : \bar{h} \in G^{k-1}\}$. The desired decomposition is:

$$S(g_1, \dots, g_{k-2}) = \begin{cases} \frac{1}{\sqrt{|G|}} \sum_{h \in G} S(g_1, \dots, g_{\frac{k-1}{2}}, h, g_{\frac{k+1}{2}}, \dots, g_{k-2}) & \text{if } k \text{ is odd} \\ S(g_1, \dots, g_{\frac{k-2}{2}}, g_{\frac{k}{2}}, g_{\frac{k}{2}}, \dots, g_{k-2}) & \text{if } k \text{ is even.} \end{cases} \quad (5)$$

(f) In what follows, we shall find it convenient to use the notation

$$\lceil x \rceil = \min\{n \in \mathbb{Z} : x \leq n\}.$$

The relations in $P(G)$ are seen to imply that for arbitrary $\bar{g}, \bar{h} \in G^{k-1}$, we have:

$$\begin{aligned}
 & S(g_1, g_2, \dots, g_{k-1})S(h_1, h_2, \dots, h_{k-1}) \\
 &= |G|^{\lceil \frac{k}{2} \rceil - 1} \left(\prod_{i=2}^{\lceil \frac{k}{2} \rceil} \delta(h_1 g_{k+1-i}, h_i) \right) \\
 &\quad \times S(h_1 g_1, h_1 g_2, \dots, h_1 g_{\lceil \frac{k}{2} \rceil}, h_{\lceil \frac{k}{2} \rceil + 1}, h_{\lceil \frac{k}{2} \rceil + 2} \dots h_{k-1})
 \end{aligned} \tag{6}$$

where $\delta(a, b)$ is zero unless $a = b$ in which case it is one.

(g) The action of Θ on $P(G)_k$ maps an orthonormal basis onto itself and consequently yields a unitary representation of Θ ; in particular, the orthogonal projection of $P(G)_k$ onto P_k^Θ is given by the usual averaging operator. So, if we define $\Theta S(\bar{g}) = \sum_{\theta \in \Theta} S(\theta(\bar{g}))$, it is seen that $\{\Theta S(\bar{g}) : [\bar{g}] \in G^k/\Theta\}$ is an orthogonal set of vectors, which clearly spans P_k^Θ (where we have written $[\bar{g}]$ to denote the orbit of \bar{g} under Θ , and G^k/Θ to denote the set of all such orbits in G^k). Finally, it is not hard to deduce from equation (6) and the relations defining $P(G)$ that if $\bar{g}, \bar{h} \in G^{k-1}$, then



$$\begin{aligned}
 & \Theta S(g_1, g_2, \dots, g_{k-1})\Theta S(h_1, h_2, \dots, h_{k-1}) \\
 &= |G|^{\lceil \frac{k}{2} \rceil - 1} \sum_{\theta'' \in \Theta} \left(\prod_{i=2}^{\lceil \frac{k}{2} \rceil} \delta(h_1 \theta''(g_{k+1-i}), h_i) \right) \\
 &\quad \times \Theta S(h_1 \theta''(g_1), h_1 \theta''(g_2), \dots, h_1 \theta''(g_{\lceil \frac{k}{2} \rceil}), h_{\lceil \frac{k}{2} \rceil + 1}, h_{\lceil \frac{k}{2} \rceil + 2} \dots h_{k-1})
 \end{aligned} \tag{7}$$

THEOREM 3.3.2. *Let G, Θ be as above, and let $G \rtimes \Theta$ denote the semi-direct product associated to this group action, and let $N = R^{G \rtimes \Theta} \subset R^\Theta = M$ denote the associated subgroup-subfactor. Then*

$$P^\Theta \cong P^{N \subset M}.$$

Proof: For notational convenience let $P_k = P_k^\Theta$, $P_{1,k} = P_{1,k}^\Theta$, $Q_k = P_k^{N \subset M}$, $Q_{1,k} = P_{1,k}^{N \subset M}$. We shall write the elements of $G \rtimes \Theta$ as ordered pairs (g, θ) with the usual multiplication $(g_1, \theta_1)(g_2, \theta_2) = (g_1 \theta_1(g_2), \theta_1 \theta_2)$. We begin by recalling some of the work in [3] which will allow us to describe

the standard invariant of P^{NCM} . We let $\boxed{q} = \frac{1}{|\Theta|} \sum_{\theta \in \Theta} \boxed{(1, \theta)}$ be the projection corresponding to the subgroup $\Theta \subset G \rtimes \Theta$. Define $B_n \in \mathcal{A}_{n,n}$ to be the following annular map in $P(G \rtimes \Theta)$:



Define the natural inclusion map

$$i : B_n(P_n) \rightarrow B_{n+1}(P_{n+1})$$

given by

$$t \mapsto B_{n+1}(t).$$

We denote this inclusion by \subset_i .

Corollary 4.5 of [3] then states that the tower

$$\begin{array}{cccc}
 Q_0 & \subset & Q_1 & \subset & Q_2 & \subset & \dots \\
 & & \cup & & & & \\
 & & Q_{1,1} & \subset & Q_{1,2} & \subset & \dots \\
 & & & & & & \dots \subset Q_n \dots \\
 & & & & & & \cup \\
 & & & & & & \dots \subset Q_{1,n} \dots
 \end{array}$$

is isomorphic to the tower:

$$\begin{aligned}
 B_0(P_0(G \rtimes \Theta)) \subset_i B_1(P_1(G \rtimes \Theta)) \subset_i B_2(P_2(G \rtimes \Theta)) \dots \\
 \cup \\
 B_1(P_{1,1}(G \rtimes \Theta)) \subset_i B_2(P_{1,2}(G \rtimes \Theta)) \dots \\
 \dots \subset_i B_n(P_n(G \rtimes \Theta)) \dots \\
 \cup \\
 \dots \subset_i B_n(P_{1,n}(G \rtimes \Theta)) \dots
 \end{aligned}$$

Using the relations in $P(G \rtimes \Theta)$ we find that

$$\begin{aligned}
 & B_n(S((g_1, \theta_1), (g_2, \theta_2), \dots, (g_{n-1}, \theta_{n-1}))) \\
 &= \frac{1}{|\Theta|^n} \sum_{\theta \in \Theta, \bar{\gamma} \in \Theta^{2n-1}} S((\theta(g_1), \gamma_1), (\theta(g_2), \gamma_2), \dots, (\theta(g_{n-1}), \gamma_{n-1}))
 \end{aligned}$$

which *only* depends on the orbit of $(g_1, g_2, \dots, g_{n-1})$ under Θ . We shall denote this sum of elements - i.e., $|\Theta|^n$ times the right side of the above equation - by $U(\bar{g})$; again, note that $U(\bar{g})$ depends only on the orbit $[\bar{g}]$ of \bar{g} under Θ . It follows then that $\{U(\bar{g})\}_{[\bar{g}] \in G^{n-1}/\Theta}$ is an orthogonal basis for $B_n(P_n(G \rtimes \Theta))$. (It is a complete set since it is the image under B_n of a basis for $P_n(G \rtimes \Theta)$. It is an orthogonal set because distinct elements are linear combinations of disjoint subsets of an orthonormal basis of $P_n(G \rtimes \Theta)$.)

Analogous to equation (6), we find - arguing this time in the group $G \rtimes \Theta$ - that if $\bar{g}, \bar{h} \in G^{k-1}$, then

$$\begin{aligned}
 & U(g_1, g_2, \dots, g_{k-1})U(h_1, h_2, \dots, h_{k-1}) \\
 &= |G|^{\lceil \frac{k}{2} \rceil - 1} |\Theta|^{k-1} \sum_{\theta'' \in \Theta} \left(\prod_{i=2}^{\lceil \frac{k}{2} \rceil} \delta(h_1 \theta''(g_{k+1-i}), h_i) \right) \\
 & \quad \times U(h_1 \theta''(g_1), h_1 \theta''(g_2), \dots, h_1 \theta''(g_{\lceil \frac{k}{2} \rceil}), h_{\lceil \frac{k}{2} \rceil + 1}, h_{\lceil \frac{k}{2} \rceil + 2} \dots h_{k-1})
 \end{aligned} \tag{8}$$

Define $\beta_k : P_k \rightarrow B_k(P_k(G \rtimes \Theta))$ by

$$\beta_k(\Theta(S(\bar{g}))) = |\Theta|^{1-k} U([\bar{g}]), \quad \bar{g} \in G^{k-1}. \tag{9}$$

By the foregoing remarks, β_k maps an orthogonal basis of P_k onto an orthogonal basis of $B_k(P_k(G \rtimes \Theta))$ and is thus a well defined bijection of vector spaces. To establish the isomorphism of towers we only need to verify that, for all k ,

1. β_k is a homomorphism,
2. $\beta_k(P_{1,k}) = B_k(P_{1,k}(G \rtimes \Theta))$,
3. $\beta_k|_{P_{k-1}} = i \circ \beta_{k-1}$.

It is a straightforward consequence of equations (6), (7) and the carefully chosen constant in Definition (9) that β_k indeed preserves multiplication and is thus a homomorphism.

The second assertion above is a consequence of Remark 3.3.1(d), and the fact that $\{U(\bar{g}) : g_1 = 1, [\bar{g}] \in G^{k-1}/\Theta\}$ is a basis for $B_k(P_{1,k}(G \rtimes \Theta))$ (the proof of which fact is analogous to that of Remark 3.3.1(d)).

Finally, the third assertion above follows from Remark 3.3.1(e). □


In the sequel, we shall economise on parentheses and write $P^{\Theta;n}$ for $(P^\Theta)^n$.

COROLLARY 3.3.3. *Suppose there exists $\bar{g}^0 \in G^{l-1}$ such that the mapping*

$$\Theta \ni \theta \mapsto \theta(\bar{g}^0) \in G^{l-1}$$

is injective. Then $P^\Theta = P^{\Theta;2l}$.

Proof: If $k \geq 2l - 1$ and $A, B \in P_k$, define $\Pi(A, B)$ to be the element of $P_{2k-2l+1}$ given by the following tangle:



It follows from the group relations that

$$\begin{aligned}
 & \Pi(T(g_1, \dots, g_{k-1}), T(h_1, \dots, h_{k-1})) \\
 &= |G|^{l-1} \left(\prod_{i=1}^{l-1} \delta(g_{k-i} h_i, 1) \right) T(g_1, \dots, g_{k-l}, h_l, \dots, h_{k-1}) . \quad (10)
 \end{aligned}$$

Let us write (as in Remark 3.3.1(g))

$$\Theta T(\bar{g}) = \sum_{\theta \in \Theta} T(\theta(\bar{g})) .$$

Then, for arbitrary $\bar{g} \in G^{2k-2l+1}$, we find, using (9), that

$$\begin{aligned}
 & \Pi(\Theta T(g_1, \dots, g_{k-l}, g_1^0, \dots, g_{l-1}^0), \Theta T((g_{l-1}^0)^{-1}, (g_{l-2}^0)^{-1}, \dots, (g_1^0)^{-1}, g_{k-l+1}, \dots, g_{2k-2l+1})) \\
 &= |G|^{l-1} \sum_{\theta, \theta'} \left(\prod_{i=1}^l \delta(\theta(g_{l-i}^0) \theta'((g_{l-i}^0)^{-1}), 1) \right) \\
 & \quad \times T(\theta(g_1), \theta(g_2), \dots, \theta(g_{k-l}), \theta'(g_{k-l+1}), \dots, \theta'(g_{2k-2l+1})) . \quad (11)
 \end{aligned}$$

Since $\theta \mapsto \theta(\bar{g}^0)$ is 1-1, it follows that

$$\prod_{i=1}^l \delta(\theta(g_{l-i}^0) \theta'((g_{l-i}^0)^{-1}), 1) = \delta(\theta(\bar{g}^0), \theta'(\bar{g}^0)) = \delta(\theta, \theta') ,$$

and thus the right side of (10) simplifies to


$$|G|^{l-1} \sum_{\theta} T(\theta(g_1), \dots, \theta(g_{2k-2l+1})) = |G|^{l-1} \Theta T(\bar{g}) .$$

Since $\{\Theta T(\bar{g})\}$ is a basis for $P_{2k-2l+1}$, we have shown $P^{\Theta; 2l} = P^{\Theta; 2l+1} = P^{\Theta; 2l+3} = P^{\Theta; 2l+7} = \dots = P^{\Theta}$.


EXAMPLE 3.3.4. We consider a few examples.

(a) Let $\Theta = \mathbb{Z}_2$ act on \mathbb{Z}_n ‘by inversion’; thus, the involutory automorphism corresponding to the non-trivial element of Θ is given by $\tau(x) = -x \forall x \in \mathbb{Z}_n$. It follows from ‘the Mackey machine’ - see [4] or [8], for instance - that the principal graph (for P^{Θ} and hence for the subgroup-subfactor corresponding to the inclusion $\mathbb{Z}_2 \subset \mathbb{Z}_n \rtimes \mathbb{Z}_2$) is given thus:

Case (i): n is odd



Case (ii): n is even



In both cases, it is clear that $P^{\ominus;1} = TL \neq P^{\ominus;2}$, while it follows from Proposition 2.2.1 that $P^{\ominus;3} = P^{\ominus;4} = P^{\ominus}$ if n is odd. (Of course, the case $n = 3$ must be discussed separately, since we get $P^{\ominus} = TL$ in this case.) Further, an argument given in [9] (see the pictorial identity within the example $G = D_{2n+1}$ towards the end) shows that $P^{\ominus;2} = P^{\ominus;3}$ for all n . In particular, the planar depth of P^{\ominus} is two, if n is odd.


On the other hand, it turns out that for even n , the planar depth of P^{\ominus} is four, as we show now. In view of the remarks of the last paragraph, it will suffice to show that $P^{\ominus;2} \neq P^{\ominus}$ in this case. Since P^{\ominus} has finite depth, it will suffice to show that $P^{\ominus;2}$ is the free product of $P(\mathbb{Z}_2)$ and $P(\mathbb{Z}_{n/2})$ (since free products necessarily have infinite depth). (See [2] for the definition and these facts about free products - or free compositions, as they are called there - of subfactors; also see [6] for free products in the planar algebra context.)

We know from Theorem 3.3.2 that $P^{\ominus} = P^{N \subset M}$, where $N = R^{\mathbb{Z}_n \rtimes \mathbb{Z}_2}$ and $M = R^{\mathbb{Z}_2}$. Then, it follows from the analysis of [3] that the free

We can deduce from Proposition 2.2.1 that $P^{\Theta;3} = P^{\Theta;4} = P^{\Theta}$. We can easily see that $P^{\Theta;1} = TL \neq P^{\Theta;2}$; we shall now proceed to show that $P^{\Theta;2} = P^{\Theta;3}$.

Let X (resp., Y) denote the sum of the 2-boxes labelled by the members of $[1] = \{1, 2, 4\}$ (resp., $[3] = \{3, 6, 5\}$). If we let Z denote the 2-box labelled by 0, then it is clear that $\{Z, X, Y\}$ is a basis for $P_2^{\Theta;2}$.


If $z, w \in G$, let us write $((z, w))$ for the element (of $P(G)_3$) obtained if we substitute z and w respectively, for E and F in the picture given by



and $[z, w] = \sum_{\theta \in \Theta} ((\theta(z), \theta(w)))$. The definitions imply - by considering the Θ -orbits in $G \times G$ - that P_3^{Θ} is linearly spanned by the set

$$\{[0, 0], [0, 1], [0, 3]\} \cup \{[1, w] : w \in G\} \cup \{[3, w] : w \in G\} .$$

Next we write (A, B, C, D) for the value of the following picture (where $A, B, C, D \in P_2^{\Theta;2}$):



and we have the following identities:

$$\begin{aligned} (Z, Z, Z, Z) &= \sqrt{7} [0, 0] \\ (Z, Z, Z, X) &= \sqrt{7} [0, 1] \\ (Z, Z, Z, Y) &= \sqrt{7} [0, 3] \end{aligned}$$

$$\begin{aligned}
 (X, Z, Y, Z) &= \sqrt{7} [1, 0] \\
 (X, X, Z, Z) &= \sqrt{7} [1, 1] \\
 (X, X, X, Z) &= \sqrt{7} [1, 2] \\
 (X, Y, X, Z) &= \sqrt{7} ([1, 3] + [1, 5]) \\
 (X, X, Y, Z) &= \sqrt{7} [1, 4] \\
 (X, Y, X, X) &= \sqrt{7} (2[1, 0] + [1, 2] + [1, 4] + [1, 5] + [1, 6]) \\
 (X, Y, Y, Z) &= \sqrt{7} [1, 6].
 \end{aligned}$$

These identities show that $[0, x], [1, x] \in P_3^{\Theta;2}$ for all $x \in G$. Similar computations show that $[3, x] \in P_3^{\Theta;2}$ for all $x \in G$. Thus, we have shown that $P_3^{\Theta} \subset P^{\Theta;2}$, so $P^{\Theta;3} \subset P^{\Theta;2}$, and $P^{\Theta;3} = P^{\Theta;2}$, as desired.

ACKNOWLEDGMENT

We would like to thank the staff of MSRI, Berkeley and the organisers of the Program on Operator Algebras for providing us a stimulating atmosphere in an idyllic ambience during the winter of 2000/01 when this work was carried out. We also wish to thank Vaughan Jones for numerous insightful conversations.

REFERENCES

1. Asaeda, M. and Haagerup, U., *Exotic subfactors of finite depth with Jones indices $(5 + \sqrt{13})/2$ and $(5 + \sqrt{17})/2$* , Comm. Math. Phys., **202**, no. 1, 1–63, 1999.
2. Bisch, D. and Jones, V., *Algebras associated to intermediate subfactors*, Invent. Math., **128**, no. 1, 89–157, 1997.
3. Bhattacharyya, Bina, and Landau, Zeph, *Intermediate standard invariants and intermediate planar algebras*, preprint.
4. Goodman, F., de la Harpe, P., and Jones, V., *Coxeter graphs and towers of algebras*, Mathematical Sciences Research Institute Publications, 14, Springer-Verlag, New York, 1989.
5. Haagerup, U., *Principal graphs of subfactors in the index range $4 < [M : N] < 3 + \sqrt{2}$* , Subfactors, (Kyuzeso, 1993), World Sci. Publishing, River Edge, NJ, 1994.
6. Jones, V.F.R., *Planar Algebras I*, New Zealand J. Math., to appear. (See www.math.berkeley.edu/~vfr)
7. Jones, V.F.R. and Sunder, V.S., *Introduction to Subfactors*, Cambridge University Press, 1997.
8. Kodiyalam, V. and Sunder, V.S., *The subgroup-subfactor*, Math. Scand., **86**, (2000), 45–74.
9. Landau, Zeph, *Exchange relation planar algebras*, Geometriae Dedicata, to appear.
10. S. Popa, *An axiomatization of the lattice of higher relative commutants of a subfactor*, Inventiones Mathematicae, vol 120, No 3, 1995.