

THE ABEL-JACOBI ISOMORPHISM ON ONE CYCLES ON THE MODULI SPACE OF VECTOR BUNDLES WITH TRIVIAL DETERMINANT ON A CURVE

JAYA NN IYER

ABSTRACT. We consider the moduli space $\mathcal{SU}_C^s(r, \mathcal{O}_C)$ of rank r stable vector bundles with trivial determinant on a smooth projective curve C of genus g . We show that the Abel-Jacobi map on the rational Chow group $CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q}$ of one cycles which are homologous to zero, is an isomorphism onto the intermediate Jacobian of the compactly supported cohomology, which is identified with the Jacobian $Jac(C) \otimes \mathbb{Q}$. This holds whenever $r \geq 2$ and $g \geq 4$.

CONTENTS

1.	Introduction	1
2.	Abel-Jacobi surjectivity for one cycles on $\mathcal{SU}_C(r, \mathcal{O}(x))$	4
3.	Hard Lefschetz isomorphism for the bottom weight odd degree rational cohomology of $\mathcal{SU}_C^s(2, \mathcal{O}_C)$	7
4.	Abel-Jacobi surjectivity for one cycles on $\mathcal{SU}_C^s(r, \mathcal{O}_C)$, $r \geq 2$	17
5.	Chow generation of one cycles on $\mathcal{SU}_C(r, \mathcal{O}_C)$	19
	References	21

1. INTRODUCTION

Suppose C is a smooth connected complex projective curve of genus g . For a line bundle L on C consider the moduli space $\mathcal{SU}_C(r, L)$ of semi-stable vector bundles of rank r and of fixed determinant L on C . In this paper we assume that either $L := \mathcal{O}_C$ or $L := \mathcal{O}_C(x)$ for some point $x \in C$. When $L = \mathcal{O}_C$ the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$ is a (singular) normal projective variety and when $L = \mathcal{O}_C(x)$ the moduli space $\mathcal{SU}_C(r, \mathcal{O}(x))$ is a smooth projective variety. The space of stable bundles $\mathcal{SU}_C^s(r, \mathcal{O}_C) \subset \mathcal{SU}_C(r, \mathcal{O}_C)$ forms a smooth quasi-projective variety. Both the moduli spaces have dimension equal to $(r^2 - 1)(g - 1)$.

⁰Mathematics Classification Number: 53C05, 53C07, 53C29

⁰Keywords: Moduli spaces, Abel-Jacobi maps, Chow groups.

It is known that these moduli spaces are unirational [Se3]. In fact, the moduli spaces $\mathcal{SU}_C(r, \mathcal{O}(-x))$ and $\mathcal{SU}_C^s(r, \mathcal{O}_C)$ are Fano manifolds [Ra], [Be]. This implies that the rational Chow group of zero cycles is trivial, i.e., $CH_0(\mathcal{SU}_C(r, L)) \otimes \mathbb{Q} \simeq \mathbb{Q}$. More generally, when X is any smooth unirational variety of dimension equal to n then it is well-known that the Hodge groups $H^p(X, \Omega_X^q) = 0$ whenever $p = 0$. In this case the intermediate Jacobian $IJ^p(X) := \frac{H^{2p-1}(X, \mathbb{C})}{F^p + H^{2p-1}(X, \mathbb{Z})}$ for $p = 2, n-1$ is an abelian variety. It is of interest to show the *weak representability* via the Abel-Jacobi map

$$CH^p(X)_{\text{hom}} \xrightarrow{AJ^p} IJ^p(X)$$

for $p = 2, n-1$ and also determine the abelian variety in terms of the geometry of X . Some examples of Fano threefolds F were shown to have *weakly representable* $CH_1(F)$ by Bloch and Murre [Bl-Mr]. When $X = \mathcal{SU}_C(2, \mathcal{O}(x))$, we have

$$\begin{aligned} CH^1(\mathcal{SU}_C(r, \mathcal{O}(x))) \otimes \mathbb{Q} &\simeq \mathbb{Q} [\text{Ra}] \\ CH^2(\mathcal{SU}_C(2, \mathcal{O}(x))) \otimes \mathbb{Q} &\simeq CH_0(C) \otimes \mathbb{Q}, \quad g = 2 \\ &\simeq CH_0(C) \otimes \mathbb{Q} \oplus \mathbb{Q}, \quad g > 2 \quad [\text{Ba-Kg-Ne}] \\ CH_1(\mathcal{SU}_C(2, \mathcal{O}(x))) \otimes \mathbb{Q} &\simeq CH_0(C) \otimes \mathbb{Q} \quad [\text{Ch-Hw}]. \end{aligned}$$

Set $n := (r^2 - 1)(g - 1)$, $AJ_1 := AJ^{n-1}$ and suppose $X = \mathcal{SU}_C(r, \mathcal{O}_C)$. Since the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$ is a singular variety, we do not have good Abel-Jacobi maps and perhaps other Chow/cohomology theories should be considered. We do not look at these theories in this paper. Instead we consider the smooth variety $\mathcal{SU}_C^s(r, \mathcal{O}_C)$ and whose cohomology groups are known to have a mixed Hodge structure. We consider the Abel-Jacobi maps which take values in the intermediate Jacobian of the (pure Hodge structure) compactly supported cohomologies. Using the results of Arapura-Sastry [Ar-Sa] in the trivial determinant case, we know that the degree 3 rational cohomology group has a pure Hodge structure of weight 3 and the intermediate Jacobian IJ^2 is identified with the Jacobian variety $Jac(C) \otimes \mathbb{Q}$. In order to study the Abel-Jacobi map AJ_1 , we firstly need to identify the appropriate target group. This is done by applying Poincaré duality to obtain an isomorphism $Jac(C) \otimes \mathbb{Q} \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C)))$.

With notations as above, we show:

Theorem 1.1. *Suppose $g \geq 3$, $r \geq 2$ and fix any point $x \in C$. Then the Abel-Jacobi map on the rational Chow group of one cycles homologous to zero*

$$(1) \quad CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{\text{hom}} \otimes \mathbb{Q} \xrightarrow{AJ_1} Jac(C) \otimes \mathbb{Q}$$

is always surjective. The same assertion is true on the non-compact smooth variety $\mathcal{SU}_C^s(r, \mathcal{O}_C)$, whenever $g \geq 4$. Furthermore, in this case, the Abel-Jacobi map extends to isomorphisms

$$\begin{aligned} CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} &\simeq Jac(C) \otimes \mathbb{Q}, \\ CH_1(\mathcal{SU}_C(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} &\simeq Jac(C) \otimes \mathbb{Q}. \end{aligned}$$

The proof of Theorem 1.1 is by combining the methods of [Ch-Hw] and the Hecke correspondence employed in [Ar-Sa]. The first assertion uses the correspondence cycle $c_2(U)$ (where U is a universal Poincaré bundle [Na-Ra]) between the Chow groups of 0-cycles on C and the codimension 2 cycles on $\mathcal{SU}_C(r, \mathcal{O}(x))$. Together with the isomorphism of the Lefschetz operator on $H^3(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$, we conclude the surjectivity of the Abel-Jacobi map on one cycles. In the case of the non compact smooth variety $\mathcal{SU}_C^s(r, \mathcal{O}_C)$, a Hecke correspondence was used in [Ar-Sa] for computing the low degree cohomology groups in terms of the cohomology of $\mathcal{SU}_C(r, \mathcal{O}(-x))$. We use this correspondence to relate the Chow group of one cycles of these two spaces, for any $r \geq 2$ and $g \geq 4$.

To conclude the theorem, we need to find a minimal generating set of one cycles on $\mathcal{SU}_C(r, \mathcal{O}_C)$. The Hecke curves [Na-Ra2] are minimal rational curves on the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$. A variant of a theorem of Kollar [Ko, Proposition 3.13.3] on Chow generation, is proved by J-M. Hwang and he observed that the Hecke curves generate the rational Chow group of one cycles on $\mathcal{SU}_C(r, \mathcal{O}_C)$, see Proposition 5.1, Corollary 5.2. This gives a surjective map

$$Jac(C) \otimes \mathbb{Q} \rightarrow CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q}.$$

Together with the Abel-Jacobi surjectivity, we conclude our main theorem.

When $r = 2$, we investigate further properties of the Chow groups and the cohomology of the moduli space $\mathcal{SU}_C(2, \mathcal{O}_C)$, in §3. We look at the codimension two cycles on the variety $\mathcal{SU}_C^s(2, \mathcal{O}_C)$ and prove the isomorphism

$$CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \simeq Jac(C) \otimes \mathbb{Q}$$

whenever $g \geq 3$, see Lemma 3.14. This isomorphism helps us in giving another proof of the surjectivity of the Abel-Jacobi map, via compatibility of the Lefschetz operator with respect to its action on the Chow groups and the cohomology groups. This proof works for $g \geq 3$ and $r = 2$.

In general, the Hard Lefschetz isomorphism is not true for the rational cohomology of open smooth varieties. In our situation, we show that the Hard Lefschetz isomorphism holds for the bottom weight cohomologies of $\mathcal{SU}_C^s(2, \mathcal{O}_C)$ in any odd degree, see Lemma 3.7. This part could be of independent interest also.

Proposition 1.2. *Suppose \mathcal{L}_s is an ample class on the moduli space $\mathcal{SU}_C^s(2, \mathcal{O}_C)$. Let $W_i H^i(\mathcal{SU}_C^s(2, \mathcal{O}_C))$ denote the bottom weight cohomology group of $\mathcal{SU}_C^s(2, \mathcal{O}_C)$, for any i . Then the Lefschetz operator \mathcal{L}_s^i gives an isomorphism*

$$\mathcal{L}_s^i : W_{n-i} H^{n-i}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q}) \xrightarrow{\sim} W_{n+i} H^{n+i}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$$

whenever $n - i$ is odd and $g \geq 3$.

This is proved using the desingularisation \mathcal{S} of $\mathcal{SU}_C(2, \mathcal{O}_C)$ constructed by Seshadri ([Se2]) and using the description of the exceptional locus. The exceptional locus has a stratification given by the rank of a conic bundle [Ba], [Ba-Se]. The computation of the low

degree cohomology of \mathcal{S} in [Ba-Se] used the Thom-Gysin sequences for the stratification. We also use similar long exact sequences of Borel-Moore homologies for the stratification of \mathcal{S} . A closer analysis reveals that the bottom weight Borel-Moore homologies $W_{-n+i}H_{n-i}$ in odd degree of the exceptional divisor are zero. This suffices to conclude the Hard Lefschetz isomorphism on the bottom weight odd degree cohomologies of $\mathcal{SU}_C^s(2, \mathcal{O}_C)$. In particular we identify the target group of AJ_1^s to be $IJ(W_{2n-3}H^{2n-3}(\mathcal{SU}_C(2, \mathcal{O}_C))) \simeq \text{Jac}(C) \otimes \mathbb{Q}$. It was pointed out by D. Arapura that at least for the degree 3 rational cohomology group, the Hard Lefschetz isomorphism can also be proved, using the Hard Lefschetz isomorphism for the intersection cohomology of $\mathcal{SU}_C(2, \mathcal{O}_C)$. This gives another way of identifying the target group of the Abel-Jacobi map AJ_1^s .

In the higher rank $r > 2$ case, we pose the question of proving the Hard Lefschetz theorem for appropriate even/odd degree cohomology groups and this could deserve some attention in the future.

Acknowledgements: This work was initiated at KIAS, Seoul, on an invitation by Jun-Muk Hwang during Feb 2009. We are grateful to him for asking this question and for all the discussions and communications we had on the subject and also for extending support and hospitality. In particular he provided the crucial results in §5. He graciously allowed this to be a single authored work, inspite of his contributions. We also thank D. Arapura for communicating to us the application of intersection cohomology. This also provided a hint to apply Poincaré duality in higher rank case.

2. ABEL-JACOBI SURJECTIVITY FOR ONE CYCLES ON $\mathcal{SU}_C(r, \mathcal{O}(x))$

Suppose C is a smooth connected projective curve defined over the complex numbers of genus g . Fix a point $x \in C$ and consider the moduli space $\mathcal{SU}_C(r, \mathcal{O}(x))$ of stable vector bundles of rank r and fixed determinant $\mathcal{O}(x)$ on C . Atiyah and Bott [At-Bo] have described the generators of the cohomology ring $H^*(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$ in terms of the characteristic classes of the Poincaré bundle. Since we are concerned only with certain cohomologies in low degree, we will recall the generators in low degrees. We also note that the moduli space $\mathcal{SU}_C(r, \mathcal{O}(x))$ is isomorphic to $\mathcal{SU}_C(r, \mathcal{O}(-x))$ given by $E \mapsto E^*$, the dual of E . In the next section we will consider the moduli space $\mathcal{SU}_C(r, \mathcal{O}(-x))$ and relate it with the results from this section.

Fix a Poincaré bundle $U \rightarrow C \times \mathcal{SU}_C(r, \mathcal{O}(x))$ and denote the i -th Chern class of U by $c_i(U) \in H^{2i}(C \times \mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$. Denote the two projections by

$$p_1 : C \times \mathcal{SU}_C(r, \mathcal{O}(x)) \rightarrow C, \quad p_2 : C \times \mathcal{SU}_C(r, \mathcal{O}(x)) \rightarrow \mathcal{SU}_C(r, \mathcal{O}(x)).$$

The cycle $c_i(U)$ acts as a correspondence between the cohomology of C and the cohomology of $\mathcal{SU}_C(r, \mathcal{O}(x))$. More precisely, there are homomorphisms:

$$\begin{aligned} H^k(C, \mathbb{Q}) &\xrightarrow{p_1^*} H^k(C \times \mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q}) \xrightarrow{\cup c_i(U)} H^{k+2i}(C \times \mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q}) \\ &\xrightarrow{p_{2*}} H^{k+2i-2}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q}). \end{aligned}$$

The composition $\Gamma_{c_i(U)} := p_{2*} \circ \cup c_i(U) \circ p_1^*$ is called the correspondence defined by the cycle $c_i(U)$.

Theorem 2.1. *The correspondence*

$$\Gamma_{c_2(U)} : H^1(C, \mathbb{Q}) \longrightarrow H^3(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$$

is an isomorphism, for $r \geq 2, g \geq 3$.

Proof. See [Na-Ra, Theorem 3] together with [Ra, Section 4 and Lemma 2.1]. The isomorphism is actually with integral coefficients. \square

Fix an ample line bundle $\mathcal{O}(1)$ on the moduli space $\mathcal{SU}_C(r, \mathcal{O}(x))$. Denote its class $H := c_1(\mathcal{O}(1)) \in H^2(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$. Set $n := \dim \mathcal{SU}_C(r, \mathcal{O}(x))$.

Corollary 2.2. *The composition*

$$H^1(C, \mathbb{Q}) \xrightarrow{\Gamma_{c_2(U)}} H^3(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q}) \xrightarrow{\cup H^{n-3}} H^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Q})$$

is an isomorphism of pure Hodge structures.

Proof. This follows from Theorem 2.1 and the Hard Lefschetz theorem. \square

Corollary 2.3. *There is an isomorphism of the intermediate Jacobians*

$$Jac(C) \longrightarrow IJ_1 := \frac{H^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{C})}{F^3 + H^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Z})}$$

induced by the composed morphism $H^{n-3} \circ \Gamma_{c_2(U)}$.

We now want to show that the isomorphisms on the intermediate Jacobians are compatible with intersections and correspondences on the rational Chow groups. More precisely, consider the classes $c_2(U) \in CH^2(\mathcal{SU}_C(\mathcal{O}(x)))$ and $H \in CH^1(\mathcal{SU}_C(r, \mathcal{O}(x)))$. Set $\Gamma_{c_i(U)}^{CH} := p_{2*} \circ \cap c_i(U) \circ p_1^*$, where

$$\begin{aligned} CH^1(C) \otimes \mathbb{Q} &\xrightarrow{p_1^*} CH^1(C \times \mathcal{SU}_C(r, \mathcal{O}(x))) \otimes \mathbb{Q} \xrightarrow{\cap c_2(U)} CH^3(C \times \mathcal{SU}_C(r, \mathcal{O}(x))) \otimes \mathbb{Q} \\ &\xrightarrow{p_{2*}} CH^2(\mathcal{SU}_C(r, \mathcal{O}(x))) \otimes \mathbb{Q}. \end{aligned}$$

We consider the correspondence

$$\Gamma_{c_2(U)}^{CH} : CH^1(C) \otimes \mathbb{Q} \longrightarrow CH^2(\mathcal{SU}_C(r, \mathcal{O}(x))) \otimes \mathbb{Q}.$$

This restricts to a correspondence on the subgroup of cycles homologous to zero:

$$\Gamma_{c_2(U)}^{CH} : CH^1(C)_{hom} \otimes \mathbb{Q} \longrightarrow CH^2(\mathcal{SU}_C(r, \mathcal{O}(x)))_{hom} \otimes \mathbb{Q}.$$

Furthermore, there is a composition of morphisms:

$$(2) \quad CH^1(C)_{hom} \otimes \mathbb{Q} \xrightarrow{\Gamma_{c_2(U)}^{CH}} CH^2(\mathcal{SU}_C(r, \mathcal{O}(x)))_{hom} \otimes \mathbb{Q} \xrightarrow{\cap H^{n-3}} CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{hom} \otimes \mathbb{Q}.$$

Set $\psi := \cap H^{n-3} \circ \Gamma_{c_2(U)}^{CH}$.

We recall the exact sequence which relates the intermediate Jacobian with the Deligne cohomology group [Es-Vi]:

$$(3) \quad 0 \longrightarrow IJ_1 \longrightarrow H_{\mathcal{D}}^{2n-2}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Z}(n-1)) \longrightarrow \text{Hodge}^{n-1} \longrightarrow 0.$$

Here $\text{Hodge}^{n-1} := \text{Image}(H_{\mathcal{D}}^{2n-2}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Z}) \rightarrow H_{\mathcal{D}}^{2n-2}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{C}) \cap H^{n-1, n-1}$.

Recall the Abel-Jacobi maps:

$$\begin{aligned} CH^1(C)_{\text{hom}} &\xrightarrow{AJ_C} Jac(C), \\ CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{\text{hom}} &\xrightarrow{AJ_1} IJ_1. \end{aligned}$$

Lemma 2.4. *The map ψ defined in (2) is compatible with the Abel-Jacobi maps AJ_C and AJ_1 . In other words, the following diagram commutes:*

$$\begin{array}{ccc} CH^1(C)_{\text{hom}} \otimes \mathbb{Q} & \xrightarrow{\psi} & CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{\text{hom}} \otimes \mathbb{Q} \\ \downarrow AJ_C & & \downarrow AJ_1 \\ Jac(C) \otimes \mathbb{Q} & \xrightarrow{\simeq} & IJ_1 \otimes \mathbb{Q}. \end{array}$$

Proof. There are cycle class maps

$$\begin{aligned} CH_1(\mathcal{SU}_C(r, \mathcal{O}(x))) &\rightarrow H_{\mathcal{D}}^{2n-2}(\mathcal{SU}_C(r, \mathcal{O}(x)), \mathbb{Z}(n-1)) \\ CH^1(C) &\rightarrow H_{\mathcal{D}}^2(C, \mathbb{Z}(1)) \end{aligned}$$

which induces the Abel-Jacobi map on the subgroups of cycles which are homologous to zero (see [Es-Vi]). In other words, using the exact sequence (3), we note that the Deligne cycle class map on the subgroup of cycles homologous to zero factors via the intermediate Jacobian and this map is the same as the Abel-Jacobi map. Furthermore, the cycle class map into the Deligne cohomology is compatible with correspondences and intersection products on the Chow groups. This implies that the above diagram in the statement of the lemma is commutative. The isomorphism on the last row of the commutative diagram is given by Corollary 2.3. \square

Corollary 2.5. *The Abel-Jacobi map*

$$AJ_1 : CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{\text{hom}} \otimes \mathbb{Q} \longrightarrow IJ_1 \otimes \mathbb{Q}$$

is surjective and is a splitting, i.e., the inverse AJ_1^{-1} is well-defined and is injective.

Proof. Use the isomorphism in the last row of the commutative diagram of Lemma 2.4 to obtain the surjectivity of AJ_1 and a splitting. \square

We know by Corollary 2.5 that the Abel-Jacobi map AJ_1 is surjective and a splitting. To show that the Abel-Jacobi map is actually an isomorphism, it suffices to show that the one-cycles on the moduli space is generated by cycles parametrised by the Jacobian $Jac(C)$. In other words there should be a surjective map

$$(4) \quad Jac(C) \otimes \mathbb{Q} \rightarrow CH_1(\mathcal{SU}_C(r, \mathcal{O}(x)))_{\text{hom}} \otimes \mathbb{Q}.$$

We will see in §5 that the assumption (4) is fulfilled for $\mathcal{SU}_C(r, \mathcal{O}_C)$, whenever $r \geq 2$ and $g \geq 4$. This will help us to conclude the desired Abel-Jacobi isomorphism.

3. HARD LEFSCHETZ ISOMORPHISM FOR THE BOTTOM WEIGHT ODD DEGREE RATIONAL COHOMOLOGY OF $\mathcal{SU}_C^s(2, \mathcal{O}_C)$

In this section, we investigate the Chow/cohomology properties of the moduli space $\mathcal{SU}_C(2, \mathcal{O}_C)$. The main result is a proof of the Hard Lefschetz isomorphism for the bottom weight odd degree rational cohomology of the moduli space $\mathcal{SU}_C(2, \mathcal{O}_C)$. Along the way, we also obtain an isomorphism of the codimension two Chow group of $\mathcal{SU}_C(2, \mathcal{O}_C)$ with the Jacobian $Jac(C)$, with \mathbb{Q} -coefficients. The Lefschetz operator acts compatibly on the Chow groups and on the cohomology groups. This enables us to give a proof of the surjectivity of the Abel-Jacobi map on one cycles, when $g \geq 3$. In the next section, we investigate the Abel-Jacobi map surjectivity for the higher rank $r \geq 2$ case, under the assumption $g \geq 4$.

The main tool is the *Hecke correspondence* which relates the Chow groups and the cohomology groups of the moduli space $\mathcal{SU}_C(2, \mathcal{O}(-x))$ with those of the moduli space $\mathcal{SU}_C(2, \mathcal{O}_C)$. We recall the Hecke correspondence used by Arapura-Sastry [Ar-Sa] to study the cohomology of the moduli space $\mathcal{SU}_C^s(r, \mathcal{O}_C)$ for any $r \geq 2$, since this correspondence will also be used in the next section.

As in the previous section, fix a point $x \in C$. Consider the moduli space $\mathcal{SU}_C(r, \mathcal{O}(-x))$ which is a smooth projective variety. Then there exist a Poincaré bundle $\mathcal{P} \rightarrow C \times \mathcal{SU}_C(r, \mathcal{O}(-x))$ and let \mathcal{P}_x denote the restriction of \mathcal{P} to $\{x\} \times \mathcal{SU}_C(r, \mathcal{O}(-x))$. Denote the projectivisation $\mathbb{P} := \mathbb{P}(\mathcal{P}_x)$ and $\pi : \mathbb{P} \rightarrow \mathcal{SU}_C(r, \mathcal{O}(-x))$ be the projection. There is a universal exact sequence on $C \times \mathbb{P}$:

$$(5) \quad 0 \rightarrow (1 \times \pi)^* \mathcal{P} \rightarrow \mathcal{V} \rightarrow \mathcal{T}_0 \rightarrow 0$$

of coherent sheaves on $C \times \mathbb{P}$ such that \mathcal{V} is a vector bundle and \mathcal{T}_0 is a sheaf supported on $\{x\} \times \mathbb{P}$, which is a line bundle on \mathbb{P} . This means that \mathbb{P} parametrises exact sequences

$$0 \rightarrow W \rightarrow V \rightarrow \mathcal{O}_Z \rightarrow 0$$

of coherent sheaves on C such that $W \in \mathcal{SU}_C(r, \mathcal{O}(-x))$ and V is a vector bundle of rank r and of trivial determinant. Here Z denotes the reduced scheme supported on $x \in C$.

It is shown in [Ar-Sa, §5] that the space \mathbb{P} is a fine moduli space of quasi-parabolic bundles and parametrises quasi-parabolic structures $V \rightarrow \mathcal{O}_Z$ whose kernel is semi-stable. In particular there is a parabolic datum Δ which attaches weights $0 < \alpha_1 < \alpha_2 < 1$ (which are assumed to be small). Then we have

Theorem 3.1. *For every parabolic stable bundle $V \rightarrow \mathcal{O}_Z$, the kernel W is semi-stable. Furthermore \mathbb{P} is the moduli space of parabolic stable (which is same as stable) bundles $\mathcal{SU}_C(r, \mathcal{O}_C, \Delta)$ and the surjection $\mathcal{V} \rightarrow \mathcal{T}_0$ is the universal family of parabolic bundles. Furthermore if $W \in \mathcal{SU}_C(r, \mathcal{O}(-x))$ then V is a semi-stable vector bundle.*

Proof. See [Ar-Sa, Theorem 5.0.3, Corollary 5.0.3]. \square

In particular there is a *Hecke diagram* relating the moduli spaces $\mathcal{SU}_C(r, \mathcal{O}(-x))$ and $\mathcal{SU}_C(r, \mathcal{O}_C)$:

$$\begin{array}{ccc} \mathbb{P} & \xrightarrow{f} & \mathcal{SU}_C(r, \mathcal{O}_C) \\ \downarrow \pi & & \\ \mathcal{SU}_C(r, \mathcal{O}(-x)). & & \end{array}$$

Furthermore, there is an open set $U := f^{-1}\mathcal{SU}_C^s(2, \mathcal{O}_C) \subset \mathbb{P}$ such that $f^{-1}V \simeq \mathbb{P}(V_x^*)$ for $V \in \mathcal{SU}_C^s(r, \mathcal{O}_C)$. In particular f restricts to a projection

$$(6) \quad f_U : U \longrightarrow \mathcal{SU}_C^s(r, \mathcal{O}_C)$$

which is a \mathbb{P}^{r-1} -bundle, see [Ar-Sa, Remark 5.0.2]. The diagram

$$(7) \quad \mathcal{SU}_C(r, \mathcal{O}(-x)) \xleftarrow{\pi} \mathbb{P} \supset U \xrightarrow{f_U} \mathcal{SU}_C^s(r, \mathcal{O}_C)$$

together with Hodge theory, projective bundle formulas and codimension estimates enabled Arapura-Sastry to compare the cohomologies of the two moduli spaces in (7) at least in low degrees. For our purpose, we recall the codimension estimate [Ar-Sa, p.17];

$$(8) \quad \text{codim}(\mathbb{P} - U) \geq 3$$

whenever $g \geq 3$. Altogether, the cohomology $H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})$ has a pure Hodge structure of weight 3 and there is an isomorphism of pure Hodge structures [Ar-Sa, Theorem 8.3.1]:

$$(9) \quad H^3(\mathcal{SU}_C(r, \mathcal{O}(-x)), \mathbb{Q}) \simeq H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}).$$

Together with the isomorphism in Theorem 2.1, there is an isomorphism of Hodge structures:

$$(10) \quad H^1(C, \mathbb{Q})(-1) \simeq H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}).$$

We would like to extend this isomorphism to the cohomology $H^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})$ as an isomorphism of Hodge structures, via the Lefschetz operator. Since $\mathcal{SU}_C^s(r, \mathcal{O}_C)$ is a non-compact smooth variety, the Hard Lefschetz theorem is not immediate. In general, this theorem does not hold for non-compact smooth varieties. Hence, we investigate the action of the Lefschetz operator on suitable sub-structures of the rational cohomology group, where the isomorphism may hold.

For this purpose, we look at the resolution \mathcal{S} of $\mathcal{SU}_C(r, \mathcal{O}_C)$ constructed by Seshadri [Se2] together with an understanding of the exceptional loci of the resolution

$$(11) \quad g : \mathcal{S} \longrightarrow \mathcal{SU}_C(r, \mathcal{O}_C).$$

This map restricts to an isomorphism $g^{-1}\mathcal{SU}_C^s(r, \mathcal{O}_C) \simeq \mathcal{SU}_C^s(r, \mathcal{O}_C)$. The variety \mathcal{S} is constructed as a moduli space of semi-stable vector bundles of rank r^2 and trivial determinant whose endomorphism algebra is a specialisation of the matrix algebra. In

general, the moduli space \mathcal{S} is a normal projective variety and it is proved to be a smooth variety only when the rank $r = 2$. Hence we assume $r = 2$ in the further discussion. See also other resolutions by Narasimhan-Ramanan [Na-Ra2] and Kirwan [Ki].

3.1. Stratification of \mathcal{S} when $r = 2$. We will recall the description of the exceptional loci of Seshadri's desingularisation given in [Ba, §3], [Ba-Se]. Recall that the singular locus of $\mathcal{SU}_C(2, \mathcal{O}_C)$ is parametrised by semi-stable bundles of the type $L \oplus L^{-1}$ for $L \in \text{Jac}(C)$. The inverse map i on $\text{Jac}(C)$ is given by $L \mapsto L^{-1}$. In other words, the Kummer variety $K(C) := \frac{\text{Jac}(C)}{\langle i \rangle}$ is precisely the singular locus. Denote the image of the set of 2^{2g} fixed points by $K(C)_{\text{fix}} \subset K(C)$. There is a stratification

$$(12) \quad \mathcal{SU}_C(2, \mathcal{O}_C) = \mathcal{SU}_C^s(2, \mathcal{O}_C) \sqcup (K(C) - K(C)_{\text{fix}}) \sqcup K(C)_{\text{fix}}.$$

The desingularisation \mathcal{S} is stratified by the rank of a natural conic bundle on \mathcal{S} [Ba, §3] and there is a filtration by closed subvarieties

$$(13) \quad \mathcal{S} = \mathcal{S}_0 \supset \mathcal{S}_1 \supset \mathcal{S}_2 \supset \mathcal{S}_3$$

such that $\mathcal{S} - \mathcal{S}_1 = g^{-1}(\mathcal{SU}_C^s(2, \mathcal{O}_C))$ and \mathcal{S}_{i+1} is the singular locus of \mathcal{S}_i .

The strata are described by the following:

Proposition 3.2. *1) The image $g(\mathcal{S}_1 - \mathcal{S}_2)$ is precisely the middle stratum. In fact $\mathcal{S}_1 - \mathcal{S}_2$ is a $\mathbb{P}^{g-2} \times \mathbb{P}^{g-2}$ bundle over $K(C) - K(C)_{\text{fix}}$.*

2) The image of \mathcal{S}_2 is precisely the deepest strata $K(C)_{\text{fix}}$ and $\mathcal{S}_2 - \mathcal{S}_3$ is the disjoint union of 2^{2g} copies of a vector bundle of rank $g - 2$ over the Grassmanian $G(2, g)$. The stratum \mathcal{S}_3 is the disjoint union of 2^{2g} copies of the Grassmanian $G(3, g)$.

Proof. See [Ba-Se, section 2]. □

We now note that the strata $\mathcal{S}_1, \mathcal{S}_2$ in the stratification $\mathcal{S} \supset \mathcal{S}_1 \supset \mathcal{S}_2 \supset \mathcal{S}_3$ are singular varieties. Moreover $\mathcal{S}_2 \subset \mathcal{S}_1$ and $\mathcal{S}_3 \subset \mathcal{S}_2$ are the singular loci respectively. Hence for our purpose, we look at the Borel-Moore homology theory (for example see [Pe-St, Chapter V, §6] for properties and the notion of weights) in the below discussion.

Before proving our main result, we recall the projective bundle formula of mixed Hodge structures:

Lemma 3.3. *Suppose $p : M \rightarrow N$ is a d -fold fibre product of \mathbb{P}^{r-1} -bundles (which need not be locally trivial in the Zariski topology) over a smooth quasi-projective variety N . Then, we have an equality of the mixed Hodge structures;*

$$W_i H^i(M, \mathbb{Q}) = \bigoplus_{j \geq 0} W_{i-2j} H^{i-2j}(N, \mathbb{Q}) \otimes H^{2j}((\mathbb{P}^{r-1})^d, \mathbb{Q}).$$

The dual statement gives the equality of the Borel-Moore homologies of M and N ;

$$W_{-i} H_i(M, \mathbb{Q}) = \bigoplus_{j \geq 0} W_{2j-i} H_{i-2j}(N, \mathbb{Q}) \otimes H_{2j}((\mathbb{P}^{r-1})^d, \mathbb{Q}).$$

Proof. The proof is basically given in [Ar-Sa, Proposition 6.3.1]. Their proof is stated in terms of the 'full' cohomology $H^i(N, \mathbb{Q})$ and $H^i(M, \mathbb{Q})$, whereas for our purpose we restrict their formula to the bottom weight cohomology of M , which gives the formula as stated. Here we use the fact that the product of projective spaces have only pure Hodge structures and that the category of \mathbb{Q} -mixed Hodge structures is semi-simple. In fact the same proof also holds for the compactly supported cohomologies $H_c^i(M, \mathbb{Q})$ and $H_c^i(N, \mathbb{Q})$ and we can restrict it on the weight i -piece, to get the formula;

$$W_i H_c^i(M, \mathbb{Q}) = \bigoplus_{j \geq 0} W_{i-2j} H_c^{i-2j}(N, \mathbb{Q}) \otimes H^{2j}((\mathbb{P}^{r-1})^d, \mathbb{Q}).$$

The dual statement follows from the definition (see [Pe-St, Definition-Lemma 6.11]) of weights on the Borel-Moore homology of a variety R ;

$$H_k(R, \mathbb{Q}) = \text{Hom}_{\mathbb{Q}}(H_c^k(R, \mathbb{Q}), \mathbb{Q}).$$

This has weights in the interval $[-k, 0]$. In particular, we have the equality $W_{-k} H_k(R, \mathbb{Q}) := \text{Hom}_{\mathbb{Q}}(W_k H_c^k(R, \mathbb{Q}), \mathbb{Q})$.

□

We also recall the following standard facts for the convenience of the reader:

Lemma 3.4. *Let X be a compact smooth variety of pure dimension n and $T \subset X$ be a closed subvariety.*

a) *Then for all k we have a nonsingular pairing of mixed Hodge structures*

$$H_T^k(X) \otimes H^{2n-k}(T) \rightarrow H_T^{2n}(X) \rightarrow \mathbb{Q}(-n).$$

Here $H_T^k(X)$ denotes the cohomology supported on T .

b) *There is a long exact sequence*

$$\rightarrow H_T^k(X) \rightarrow H^k(X) \rightarrow H^k(X - T) \rightarrow .$$

The group $H_T^k(X)$ has only weights $\geq k$ and the image of this group in $H^k(X)$ has weight k . In other words, the image is identified with the image of the group $W_k H_T^k(X)$.

c) *(Poincaré duality) There is an isomorphism*

$$H_T^{2n-k}(X) \simeq H_k(T)(-n)$$

of \mathbb{Q} -mixed Hodge structures.

Proof. a) See the discussion above [Pe-St, Corollary 6.14].

b) See [Pe-St, Corollary 6.14] and noting that $H^k(X)$ has a pure Hodge structure of weight k .

c) See [Ja, p.82, h) and p.92, Example 6.9].

□

Corollary 3.5. *The weight $(-i)$ piece $W_{-i} H_i(\mathcal{S}_1, \mathbb{Q})$ of the homology $H_i(\mathcal{S}_1, \mathbb{Q})$ of \mathcal{S}_1 is zero, whenever i is odd.*

Proof. We firstly note that the involution i acts as (-1) on the odd homology (respectively $(+1)$ on the even homology) of the Jacobian $J(C)$. The i -fixed homology classes precisely correspond to the homology classes on the quotient variety $K(C)$. This means that the homology of the Kummer variety $K(C)$ is zero in odd degrees. Look at the long exact homology sequence for the triple $(K(C), K(C) - K(C)_{fix}, K(C)_{fix})$ (see [Ja, p.81, §6, f]):

$$\rightarrow H_i(K(C)_{fix}, \mathbb{Q}) \rightarrow H_i(K(C), \mathbb{Q}) \rightarrow H_i(K(C) - K(C)_{fix}, \mathbb{Q}) \rightarrow .$$

Since $K(C)_{fix}$ is a finite set of points, we conclude that the homology of the open variety $K(C) - K(C)_{fix}$ is also zero in odd degrees, except in degree $i = 1$. In this case, we obtain an injectivity $H_1(K(C) - K(C)_{fix}, \mathbb{Q}) \hookrightarrow H_0(K(C)_{fix}, \mathbb{Q})$. Since $H_0(K(C)_{fix}, \mathbb{Q})$ has weight 0, we conclude that $H_1(K(C) - K(C)_{fix}, \mathbb{Q})$ also has only weight 0. In particular, we have the vanishing $W_{-1}H_1(K(C) - K(C)_{fix}, \mathbb{Q}) = 0$.

Now we look at the homology of the triple $(\mathcal{S}_2, \mathcal{S}_2 - \mathcal{S}_3, \mathcal{S}_3)$: the long exact homology sequence for this triple is

$$\rightarrow H_i(\mathcal{S}_3, \mathbb{Q}) \rightarrow H_i(\mathcal{S}_2, \mathbb{Q}) \rightarrow H_i(\mathcal{S}_2 - \mathcal{S}_3, \mathbb{Q}) \rightarrow .$$

Note that the Grassmanian and vector bundles over Grassmanians have only algebraic homology, i.e., have homology only in even degrees. By Proposition 3.2 2), we know that $\mathcal{S}_2 - \mathcal{S}_3$ and \mathcal{S}_3 is made of such objects. Hence we conclude that \mathcal{S}_2 has all of its odd degree cohomology as zero.

Now look at the long exact homology sequence for the triple $(\mathcal{S}_1, \mathcal{S}_1 - \mathcal{S}_2, \mathcal{S}_2)$: here we note that the homology of the projective bundles or more generally flag varieties over a variety are generated by the homology of the base variety and the standard homology classes of powers of $\mathcal{O}(1)$ on these bundles. In particular these standard classes contribute only in the even homology of the total space. Since we have noticed above that $K(C) - K(C)_{fix}$ has vanishing odd degree homology in degrees > 1 , and in degree one case we have $W_{-1}H_1(K(C) - K(C)_{fix}, \mathbb{Q}) = 0$, it follows that $\mathcal{S}_1 - \mathcal{S}_2$ has in odd degree i , the vanishing of the bottom weight cohomology $W_{-i}H_i(\mathcal{S}_1 - \mathcal{S}_2, \mathbb{Q}) = 0$, by Lemma 3.3. We use the long exact homology sequence for the triple $(\mathcal{S}_1, \mathcal{S}_1 - \mathcal{S}_2, \mathcal{S}_2)$ and since \mathcal{S}_2 has vanishing odd degree homology we conclude, for an odd degree i , the vanishing of the bottom weight cohomology $W_{-i}H^i(\mathcal{S}_1, \mathbb{Q}) = 0$. \square

3.2. Hard Lefschetz isomorphism for $W_{odd}H^{odd}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$. Fix an ample class \mathcal{L} on the moduli space \mathcal{S} which restricts to an ample class \mathcal{L}_s on $\mathcal{SU}_C^s(2, \mathcal{O}_C)$. Let $n := \dim \mathcal{S}$. Now we look at the long exact cohomology sequence for the triple $(\mathcal{S}, \mathcal{SU}_C^s(2, \mathcal{O}_C), \mathcal{S}_1)$ which is compatible with the Lefschetz operators:

$$\begin{array}{ccccc} \rightarrow H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{g} & H^{n-i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{h} & H^{n-i}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q}) \rightarrow \\ & & \downarrow \cup \mathcal{L}^i & & \downarrow \cup \mathcal{L}_s^i \\ \rightarrow H_{\mathcal{S}_1}^{n+i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{g'} & H^{n+i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{h'} & H^{n+i}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q}) \rightarrow . \end{array}$$

Here $H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q})$ and $H_{\mathcal{S}_1}^{n+i}(\mathcal{S}, \mathbb{Q})$ denotes the cohomology supported on \mathcal{S}_1 .

Lemma 3.6. *The image of the cohomology $H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q})$ (respectively $H_{\mathcal{S}_1}^{n+i}(\mathcal{S}, \mathbb{Q})$) under g (respectively g') is zero, whenever $n - i$ is odd. In particular the maps h, h' in the above long exact sequences are injective whenever $n - i$ is odd.*

Proof. Since $\mathcal{S}_1 \subset \mathcal{S}$ is a divisor, we can identify the group $H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q})$ with the Borel–Moore homology $H_{n+i}(\mathcal{S}_1, \mathbb{Q}(n))$, which is actually the group $H_{n+i}(\mathcal{S}_1, \mathbb{Q})(-n)$, see Lemma 3.4 c). This group has weights $2(n) - (n + i) = n - i$ and higher, and the image of $H_{n+i}(\mathcal{S}_1, \mathbb{Q}(n))$ coincides with the image of $W_{n-i}(H_{n+i}(\mathcal{S}_1, \mathbb{Q})(-n))$, see Lemma 3.4 b).

Now we notice that since $\mathbb{Q}(-n)$ has weight $(2n)$, the group $W_{n-i}(H_{n+i}(\mathcal{S}_1, \mathbb{Q})(-n))$ is identified with the group $(W_{-n-i}H_{n+i}(\mathcal{S}_1, \mathbb{Q}))(-n)$. Since $n - i, n + i$ are odd, by Corollary 3.5, we know that $W_{-n-i}H_{n+i}(\mathcal{S}_1, \mathbb{Q}) = 0$. Hence we conclude that the image of the cohomology $H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q})$ under g is zero. A similar computation shows that the group $H_{\mathcal{S}_1}^{n+i}(\mathcal{S}, \mathbb{Q})$ is identified with the group $W_{-n+i}H_{n-i}(\mathcal{S}_1, \mathbb{Q})(-n)$ and hence its image under g' is also zero. \square

Since the images of h and h' correspond to the bottom weight cohomology of $\mathcal{SU}_C^s(2, \mathcal{O}_C)$ (see [De, p.39, Corollaire 3.2.17]), we can rewrite the above sequences as the following commutative diagram of exact sequences;

$$\begin{array}{ccccccc} \rightarrow H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{g} & H^{n-i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{h} & W_{n-i}H^{n-i}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}) & \rightarrow & 0 \\ & & \downarrow \cup \mathcal{L}_{\mathcal{S}_1}^i & & \downarrow \cup \mathcal{L}^i & & \downarrow \cup (\mathcal{L}_s^i)_W \\ \rightarrow H_{\mathcal{S}_1}^{n+i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{g'} & H^{n+i}(\mathcal{S}, \mathbb{Q}) & \xrightarrow{h'} & W_{n+i}H^{n+i}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}) & \rightarrow & 0. \end{array}$$

Lemma 3.7. *The map $\cup (\mathcal{L}_s^i)_W$ between the bottom weight cohomologies in the above long exact sequences is an isomorphism, whenever $n - i$ is odd, as a morphism of Hodge structures.*

Proof. Since the line bundle \mathcal{L} restricts to an ample class on $\mathcal{SU}_C^s(2, \mathcal{O}_C)$, the Lefschetz operator $\cup \mathcal{L}^i$ induces an operator on the cohomology $H_{\mathcal{S}_1}^{n-i}(\mathcal{S}, \mathbb{Q})$. Applying the Hard Lefschetz isomorphism on \mathcal{S} , we have the isomorphism of the operator $\cup \mathcal{L}^i$, as morphisms of Hodge structures. By Lemma 3.6, we deduce that the morphisms h and h' are injective and onto the bottom weight cohomologies. Since $\cup \mathcal{L}^i$ is an isomorphism we obtain that $\cup (\mathcal{L}_s^i)_W$ is also an isomorphism. \square

Lemma 3.8. *The Lefschetz operator*

$$\cup \mathcal{L}_s^{n-3} : H^3(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q}) \rightarrow H^{2n-3}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$$

is injective as a morphism of Hodge structures.

Proof. Consider the above long exact sequences when $n - i = 3, n + i = 2n - 3$. Using Lemma 3.7, we know that the maps h, h' in the above long exact cohomology sequence on the cohomology are injective. Since the Hard Lefschetz isomorphism holds on \mathcal{S} , the operator $\cup \mathcal{L}^{n-3}$ is an isomorphism. Furthermore, by [Ar-Sa, Theorem 8.3.1], the

cohomology $H^3(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$ has a pure Hodge structure and hence the map h is an isomorphism onto $H^3(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$. This implies that $\cup \mathcal{L}_s^{n-3}$ is injective. Moreover, the maps in the above long exact sequence are morphisms of mixed Hodge structures and since $\cup \mathcal{L}^{n-3}$ is an isomorphism of Hodge structure, we obtain the last assertion. \square

Corollary 3.9. *There is an injectivity*

$$H^1(C, \mathbb{Q}) \hookrightarrow H^{2n-3}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$$

of Hodge structures. In particular the intermediate Jacobian

$$IJ_1(W_{2n-3}) \otimes \mathbb{Q} := \frac{W_{2n-3}H^{2n-3}(\mathcal{SU}_C^s(2, \mathcal{O}_C, \mathbb{C}))}{F^{n-1} + W_{2n-3}H^{2n-3}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})}$$

is isomorphic to the Jacobian $Jac(C) \otimes \mathbb{Q}$.

Proof. Use (10) together with Lemma 3.8 to conclude that $H^1(C, \mathbb{Q})$ is a sub-Hodge structure of the mixed Hodge structure on $H^{2n-3}(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q})$. Using Lemma 3.7, we deduce the last assertion. \square

It was communicated to us by D. Arapura [Ar] that it may be possible to apply the Hard Lefschetz isomorphism on the intersection cohomology of the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$, at least in low degrees, to get the desired isomorphism. Indeed, we check this in the following discussion. We firstly recall some facts/notations on the intersection (co)homology [Go-Ma], [Go-Ma2]. Suppose X is a complex algebraic variety which is not necessarily compact and $l := \dim X$. Let $Z \subset X$ be a closed subvariety of pure dimension d and $U := X - Z$ be the open complementary subset of X . Denote

$$\begin{aligned} IH_c^k(X) &= IH_{2l-k}^k(X) \\ IH^k(X) &= IH_{2l-k}^{BM}(X). \end{aligned}$$

Here $IH_k(X)$ denotes the middle-perversity intersection homology group of X and $IH_k^{BM}(X)$ denotes the middle-perversity Borel-Moore homology group of X .

Lemma 3.10. *Let X be a compact algebraic variety of dimension l and $Z \subset X$ is a closed subvariety of dimension d with $U := X - Z$. Then with \mathbb{Q} -coefficients, we have*

$$\begin{aligned} IH^k(X) &\simeq H^k(U) \text{ for } k < l - d \\ &\simeq H_c^k(U) \text{ for } k > l - d. \end{aligned}$$

These are isomorphisms of \mathbb{Q} -mixed Hodge structures.

Proof. See [Du, p.991, Proposition 3]. \square

We also recall the famous theorem of Beilinson-Bernstein-Deligne-Gabber:

Theorem 3.11. [BBDe] *The Hard Lefschetz isomorphism holds for the intersection cohomology.*

This can be applied to our moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$ to obtain the Hard Lefschetz isomorphism at least in the degree 3 rational cohomology group;

Corollary 3.12. *Suppose \mathcal{L} is an ample class on the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$, for $r \geq 3$ and $g \geq 4$. Then there is a Hard Lefschetz isomorphism*

$$H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}) \stackrel{\cup \mathcal{L}^{n-3}}{\simeq} H_c^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C), \mathbb{Q}).$$

In particular, we have an isomorphism of the intermediate Jacobians:

$$Jac(C) \otimes \mathbb{Q} \simeq IJ(H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})) \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C), \mathbb{Q})).$$

Proof. By Theorem 3.11, we have an isomorphism of the intersection cohomology groups of $\mathcal{SU}_C(r, \mathcal{O}_C)$:

$$IH^3(\mathcal{SU}_C(r, \mathcal{O}_C)) \stackrel{\cup \mathcal{L}^{n-3}}{\simeq} IH^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C))$$

given by the Lefschetz operator. Recall the codimension estimate [Ar-Sa, Remark 5.1.1]:

$$\text{codim}(\mathcal{SU}_C(r, \mathcal{O}_C) - \mathcal{SU}_C^s(r, \mathcal{O}_C)) > 5$$

whenever $r \geq 3$ and $g \geq 4$. Then, by Lemma 3.10, we have the isomorphisms

$$IH^3(\mathcal{SU}_C(r, \mathcal{O}_C)) \simeq H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})$$

and

$$IH^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C)) \simeq H_c^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C), \mathbb{Q}).$$

This gives the Hard Lefschetz isomorphism in the degree 3 rational cohomology group of $\mathcal{SU}_C(r, \mathcal{O}_C)$. The last claim follows because all the above isomorphisms are isomorphisms of \mathbb{Q} -mixed Hodge structures and using the identification in (10). \square

In the following subsection, we show the surjectivity of the Abel-Jacobi map when $r = 2$ and $g \geq 3$. We will firstly determine the codimension two rational Chow group $CH^2(\mathcal{SU}_C(2, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q}$. Then we will apply the Hard Lefschetz isomorphism proved above and the compatibility of the Lefschetz operator acting on the Chow groups and the cohomology. In §4, we will show that the Abel-Jacobi map is surjective for any $r \geq 2$ and $g \geq 4$, by another method.

3.3. Surjectivity of the Abel–Jacobi map AJ_1^s onto $Jac(C)$, $r = 2$. Consider the Abel–Jacobi map

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} \rightarrow IJ_1(W_{2n-3}) \otimes \mathbb{Q} \simeq Jac(C) \otimes \mathbb{Q}.$$

using the identifications in Corollary 3.9 or Corollary 3.12.

We will prove that the map AJ_1^s is surjective.

Recall the Hecke diagram from the previous subsection:

$$\begin{array}{ccc} \mathbb{P} & \xrightarrow{f} & \mathcal{SU}_C(2, \mathcal{O}_C) \\ \downarrow \pi & & \\ \mathcal{SU}_C(2, \mathcal{O}(-x)). & & \end{array}$$

Denote the intermediate Jacobians of a smooth projective variety Y of dimension l by

$$IJ^2(Y) := \frac{H^3(Y, \mathbb{C})}{F^2 + H^3(Y, \mathbb{Z})}, \quad IJ_1(Y) := \frac{H^{2l-3}(Y, \mathbb{C})}{F^l + H^{2l-3}(Y, \mathbb{Z})}.$$

We will use these notations for the spaces $\mathcal{SU}_C(2, \mathcal{O}(-x))$ and the \mathbb{P}^1 -bundle \mathbb{P} .

Lemma 3.13. *a) We have the following isomorphisms of the intermediate Jacobians*

$$\begin{aligned} IJ_1(\mathbb{P}) &\simeq IJ_1(\mathcal{SU}_C(2, \mathcal{O}(-x))) \\ IJ^2(\mathcal{SU}_C(2, \mathcal{O}(-x))) &\simeq Jac(C) \otimes \mathbb{Q}. \end{aligned}$$

b) Similar isomorphisms hold for the Chow groups:

$$\begin{aligned} CH_1(\mathbb{P})_{hom} &\simeq \alpha.\pi^*CH_1(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom}, \\ CH^2(\mathbb{P})_{hom} &\simeq CH^2(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom}, \\ CH^2(U)_{hom} \otimes \mathbb{Q} &\xleftarrow{\simeq} CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q}. \end{aligned}$$

Here $\alpha := c_1(\mathcal{O}_{\mathbb{P}}(1))$.

Proof. a) There is a decomposition

$$H^{2n-1}(\mathbb{P}, \mathbb{Q}) = \pi^*H^{2n-1}(\mathcal{SU}_C(2, \mathcal{O}(-x)), \mathbb{Q}) \oplus \alpha.\pi^*H^{2n-3}(\mathcal{SU}_C(2, \mathcal{O}(-x)), \mathbb{Q}).$$

The first cohomology piece is zero since it is dual to $H^1(\mathcal{SU}_C(2, \mathcal{O}(-x)), \mathbb{Q})$ which is zero. This implies that there is an isomorphism of the intermediate Jacobians

$$IJ_1(\mathcal{SU}_C(2, \mathcal{O}(-x))) \simeq IJ_1(\mathbb{P}).$$

Again, by Hard Lefschetz isomorphism, we have the isomorphism

$$IJ^2(\mathcal{SU}_C(2, \mathcal{O}(-x))) \simeq Jac(C) \otimes \mathbb{Q}.$$

b) Using the projective bundle formula [Fu, p.64], similar statements hold on the Chow groups of cycles homologous to zero, as asserted. Indeed, we note that

$$\begin{aligned} CH^2(\mathbb{P}) \otimes \mathbb{Q} &= CH_{n-1}(\mathbb{P}) \otimes \mathbb{Q} \\ &= \pi^*CH_{n-2}(\mathcal{SU}_C(2, \mathcal{O}(-x))) \otimes \mathbb{Q} + \alpha.\pi^*CH_{n-1}(\mathcal{SU}_C(2, \mathcal{O}(-x))) \otimes \mathbb{Q}. \end{aligned}$$

Since $CH_{n-1}(\mathcal{SU}_C(2, \mathcal{O}(-x))) \otimes \mathbb{Q} = \mathbb{Q}$ ([Ra, Proposition 3.4]), we conclude the equality

$$CH^2(\mathbb{P})_{hom} \otimes \mathbb{Q} = CH^2(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom} \otimes \mathbb{Q}.$$

Also, since $CH_0(\mathcal{SU}_C(2, \mathcal{O}(-x))) \otimes \mathbb{Q} = \mathbb{Q}$, the projective bundle formula gives the equality

$$CH_1(\mathbb{P})_{hom} \otimes \mathbb{Q} = \alpha.\pi^*CH_1(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom} \otimes \mathbb{Q}.$$

Similarly, since $\text{Pic}(\mathcal{SU}_C^s(2, \mathcal{O}_C)) \otimes \mathbb{Q} = \mathbb{Q}$, we conclude the isomorphism

$$CH^2(U)_{hom} \otimes \mathbb{Q} \xleftarrow{\simeq} CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q}.$$

□

Lemma 3.14. *There is a commutative diagram*

$$\begin{array}{ccc} CH^2(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom} \otimes \mathbb{Q} & \xrightarrow{\simeq} & CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \\ \downarrow AJ^2 & & \downarrow AJ_s^2 \\ Jac(C) \otimes \mathbb{Q} & \xrightarrow{\simeq} & Jac(C) \otimes \mathbb{Q}. \end{array}$$

Furthermore, the maps AJ^2 and AJ_s^2 are isomorphisms, whenever $g \geq 3$.

Proof. Firstly, by Lemma 3.13, we have an isomorphism $IJ^2(\mathcal{SU}_C(2, \mathcal{O}(-x))) \simeq Jac(C) \otimes \mathbb{Q}$ and by (10) there is an isomorphism $Jac(C) \otimes \mathbb{Q} \simeq IJ^2(\mathcal{SU}_C^s(2, \mathcal{O}_C)) \otimes \mathbb{Q}$. Hence the maps AJ^2 and AJ_s^2 are just the Abel-Jacobi maps. Consider the following maps on the cohomology

$$(14) \quad H^3(\mathcal{SU}_C(2, \mathcal{O}(-x)), \mathbb{Q}) \xrightarrow{\pi^*} H^3(\mathbb{P}, \mathbb{Q}) \xrightarrow{s^0} H^3(U, \mathbb{Q}) \xleftarrow{f_U^*} H^3(\mathcal{SU}_C^s(2, \mathcal{O}_C), \mathbb{Q}).$$

Then all the above maps are isomorphisms, by [Ar-Sa, Theorem 8.3.1]. These are also isomorphisms of pure Hodge structures and hence induce isomorphisms of the associated intermediate Jacobians

$$(15) \quad IJ^2(\mathcal{SU}_C(2, \mathcal{O}(-x))) \simeq IJ^2(\mathbb{P}) \simeq IJ^2(U) \simeq IJ^2(\mathcal{SU}_C^s(2, \mathcal{O}_C)).$$

Similarly there are maps between the rational Chow groups

$$(16) \quad CH^2(\mathcal{SU}_C(2, \mathcal{O}(-x)))_{hom} \otimes \mathbb{Q} \xrightarrow{\pi^*} CH^2(\mathbb{P})_{hom} \otimes \mathbb{Q} \xrightarrow{t^0} CH^2(U)_{hom} \otimes \mathbb{Q} \xleftarrow{f_U^*} CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q}.$$

Using Lemma 3.13, we conclude that π^* and f_U^* are isomorphisms. Also, there is a localization exact sequence (see [Fu, Proposition 1.8, p.21])

$$\rightarrow CH_{n-1}(\mathbb{P} - U) \otimes \mathbb{Q} \rightarrow CH_{n-1}(\mathbb{P}) \otimes \mathbb{Q} \xrightarrow{t^0} CH_{n-1}(U) \otimes \mathbb{Q} \rightarrow 0.$$

Using the codimension estimate (8), we conclude that $CH_{n-1}(\mathbb{P} - U) \otimes \mathbb{Q} = 0$, i.e., t^0 is an isomorphism, whenever $g \geq 3$. This implies that all the maps in (16) are isomorphisms and compatible with the Abel-Jacobi maps into the objects in (15). This gives the commutative diagram as in the statement of the lemma. By [Ba-Kg-Ne, p.10], we know that AJ^2 is an isomorphism. This implies that AJ_s^2 is also an isomorphism. □

Lemma 3.15. *There is a commutative diagram*

$$\begin{array}{ccc} CH^2(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} & \xrightarrow{\mathcal{L}^{n-3}} & CH_1(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \\ \downarrow AJ_s^2 & & \downarrow AJ_1^s \\ Jac(C) \otimes \mathbb{Q} & \xrightarrow{\mathcal{L}_s^{n-3}} & IJ_1(W_{2n-3}) \otimes \mathbb{Q}. \end{array}$$

such that \mathcal{L}_s^{n-3} maps isomorphically onto $Jac(C) \otimes \mathbb{Q} \simeq IJ_1(W_{2n-3}) \otimes \mathbb{Q}$.

Proof. We just need to note that the Lefschetz operator acts compatibly on the Chow groups and on the cohomology as a morphism of Hodge structures. This gives the commutative diagram. Also, by Lemma 3.8 and Corollary 3.9, the map \mathcal{L}_s^{n-3} maps isomorphically onto $Jac(C) \otimes \mathbb{Q}$. \square

Consider the Abel–Jacobi map

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \xrightarrow{AJ_1^s} IJ_1(W_{2n-3}) \otimes \mathbb{Q} \simeq Jac(C) \otimes \mathbb{Q}.$$

Corollary 3.16. *The Abel–Jacobi map AJ_1^s is surjective.*

Proof. We use the commutative diagram in Lemma 3.15 and the isomorphism of \mathcal{L}_s^{n-3} onto $Jac(C) \otimes \mathbb{Q}$. By Lemma 3.14, we know that the map AJ_s^2 is an isomorphism. Hence we conclude that the map AJ_1^s is surjective. \square

4. ABEL-JACOBI SURJECTIVITY FOR ONE CYCLES ON $\mathcal{SU}_C^s(r, \mathcal{O}_C)$, $r \geq 2$

In this section, we investigate the Abel–Jacobi map

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \rightarrow IJ \otimes \mathbb{Q}$$

for any rank $r \geq 2$. The first problem is to find an appropriate identification of the group $IJ \otimes \mathbb{Q}$ and the second problem is to prove the surjectivity of this map. When $r = 2$ and $g \geq 3$, the answer to the first problem was given by the Hard Lefschetz isomorphism (proved via explicit computations which was possible only for the case $r = 2$). The answer to the second problem essentially required us to prove the isomorphism $CH^2(\mathcal{SU}_C(2, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \simeq Jac(C) \otimes \mathbb{Q}$ (which we know only in the case $r = 2$) and use the compatibility of the Lefschetz operator. When $r \geq 2$ and $g \geq 4$, we will see below that the answer to the above problems can be given via an application of the Poincaré duality and using the projective bundle formulas.

Now we consider the Abel–Jacobi map;

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \rightarrow IJ(H_c^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}_C), \mathbb{Q})).$$

As shown in the $r = 2$ case (see Lemma 2.3), we have the following isomorphisms in the higher rank $r \geq 2$ and $g \geq 4$ case as well:

Lemma 4.1. *a) We have the following isomorphisms of the intermediate Jacobians*

- 1) $IJ(H^{2n-1}(\mathbb{P}, \mathbb{Q})) \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(-x)), \mathbb{Q})),$
- 2) $IJ(H_c^{2n-1}(U, \mathbb{Q})) \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})),$
- 3) $IJ(H^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(-x)), \mathbb{Q})) \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}))$
- 4) $Jac(C) \otimes \mathbb{Q} \simeq IJ(H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})).$

b) Similar isomorphisms hold for the Chow groups:

$$\begin{aligned} CH_1(\mathbb{P})_{hom} &\simeq \alpha.\pi^*CH_1(\mathcal{SU}_C(r, \mathcal{O}(-x)))_{hom}, \\ CH_1(U)_{hom} &\simeq \beta.f_U^*CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{hom}. \end{aligned}$$

Here $\alpha := c_1(\mathcal{O}_{\mathbb{P}}(1))$ and $\beta := c_1(\mathcal{O}_{U/\mathcal{SU}_C^s(r, \mathcal{O}_C)}(1))$.

Proof. a) The proof of 1) is the same as in Lemma 2.3 a). Next for 2), we have a commutative diagram

$$\begin{array}{ccc} H^3(U, \mathbb{Q}) & \xrightarrow{f_U^*} & H^3(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}) \\ \downarrow \simeq & & \downarrow \simeq \\ H_c^{2n-1}(U, \mathbb{Q}) & \xleftarrow{\beta.f_U^*} & H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q}). \end{array}$$

The vertical isomorphisms are by Poincaré duality (we have omitted the Tate twists here). This implies that the map $\beta.f_U^*$ is an isomorphism of pure Hodge structures (since $H^3(U, \mathbb{Q})$ has a pure Hodge structure). The isomorphism in 3) is obtained from the commutative diagram:

$$\begin{array}{ccc} H^3(\mathbb{P}, \mathbb{Q}) & \xrightarrow{j'} & H^3(U, \mathbb{Q}) \\ \downarrow \simeq & & \downarrow \simeq \\ H^{2n-1}(\mathbb{P}, \mathbb{Q}) & \xrightarrow{j} & H_c^{2n-1}(U, \mathbb{Q}). \end{array}$$

The vertical isomorphisms are by Poincaré duality and the isomorphism of j' is due to the codimension estimate in [Ar-Sa, p.17]. This shows that j is an isomorphism of pure Hodge structures. Combining the isomorphisms of 1) and 2), we obtain 3) and 4).

b) The proof is the same as in Lemma 2.3 b). \square

Corollary 4.2. *There is a commutative diagram of the Abel-Jacobi maps, for $r \geq 3$, $g \geq 4$;*

$$\begin{array}{ccccc} CH_1(\mathbb{P})_{hom} \otimes \mathbb{Q} & \xrightarrow{h} & CH_1(U)_{hom} \otimes \mathbb{Q} & \simeq & CH_1(\mathcal{SU}_C(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \\ \downarrow AJ_1^{\mathbb{P}} & & \downarrow AJ_1^U & & \downarrow AJ_1^s \\ IJ(H^{2n-1}(\mathbb{P}, \mathbb{Q})) & \xrightarrow{\simeq} & IJ(H_c^{2n-1}(U, \mathbb{Q})) & \simeq & IJ(H_c^{2n-3}(\mathcal{SU}_C^s(r, \mathcal{O}_C), \mathbb{Q})). \end{array}$$

Moreover the Abel-Jacobi map AJ_1^s is surjective. Here h denotes the restriction of the one cycles on U .

Proof. The assertion on the commutativity of the above diagram basically follows from the isomorphisms in Lemma 4.1. Since $AJ_1^{\mathbb{P}}$ is the same as $AJ_1 : CH_1(\mathcal{SU}_C(r, \mathcal{O}(-x)))_{hom} \otimes \mathbb{Q} \rightarrow IJ(H^{2n-3}(\mathcal{SU}_C(r, \mathcal{O}(-x))))$, it follows from Corollary 2.5 that the map $AJ_1^{\mathbb{P}}$ is surjective. This implies that the map AJ_1^s is also surjective. \square

5. CHOW GENERATION OF ONE CYCLES ON $\mathcal{SU}_C(r, \mathcal{O}_C)$

A study of Fano manifolds with Picard number one with respect to the geometry of the variety of tangent directions to the minimal rational curves, has been studied by J-M. Hwang and N. Mok in a series of papers (see [Hw-Mo] for a survey). The moduli space $\mathcal{SU}_C^s(2, \mathcal{O}_C)$ is a Fano manifold with Picard number one and the minimal rational curves are the 'Hecke curves' introduced by Ramanan and Narasimhan [Na-Ra2]. Suppose \mathcal{L} is the ample generator of $\mathcal{SU}_C(2, \mathcal{O}_C)$ then the dualizing class K is equal to $-4\mathcal{L}$ ([Be]). Furthermore, a Hecke curve has degree 4 with respect to $-K$. Hence it has degree one with respect to \mathcal{L} (see also [Hw]). In this section, we include the results of J-M. Hwang which shows that the Hecke curves generate the Chow group of one cycles on the moduli space $\mathcal{SU}_C(2, \mathcal{O}_C)$.

With notations as in the previous section or [Ar-Sa], there is a fibration

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f} & \mathcal{SU}_C(r, \mathcal{O}_C) \\ \downarrow \pi & & \\ \mathcal{S} & & \\ \downarrow \psi & & \\ C & & \end{array}$$

such that the fibre of the morphism ψ at a point $x \in C$ is the moduli space $\mathcal{SU}_C(r, \mathcal{O}(-x))$. The variety \mathcal{P} is a \mathbb{P}^{r-1} -bundle over \mathcal{S} and restricting over a fibre $\psi^{-1}(x)$ gives precisely the Hecke correspondence used in the previous section. The image under f of the lines in the fibres of the projection π are the Hecke curves on $\mathcal{SU}_C(r, \mathcal{O}_C)$.

Firstly, we look at a variant of a theorem of Kollar [Ko, Proposition 3.13.3], on the Chow generation of one cycles on a variety, due to H-M. Hwang:

Proposition 5.1. (*Hwang*) *Let X be a normal projective variety. Let $\mathcal{M} \subset \text{Chow}(X)$ be a closed subscheme of the Chow scheme of X such that all members of \mathcal{M} are irreducible and reduced curves on X . For a general point $x \in X$, let $\mathcal{C}_x \subset \mathbb{P}T_x(X)$ be the closure of the union of the tangent vectors to members of \mathcal{M} passing through x , which are smooth at x . Suppose that \mathcal{C}_x is non-degenerate in $\mathbb{P}T_x(X)$. Then the Chow group of 1-cycles $\text{CH}_1(X) \otimes \mathbb{Q}$ is generated by members of \mathcal{M} .*

Proof. Applying Theorem IV.4.13 of [Ko], we have an open subvariety $X^\circ \subset X$ and a morphism $\pi : X^\circ \rightarrow Z^\circ$ with connected fibers such that for every $z \in Z^\circ$, any two general points of $\pi^{-1}(z)$ can be connected by a connected chain consisting of members of \mathcal{M} of length at most $\dim X$ and all members of \mathcal{M} through a point $x \in \pi^{-1}(z)$ are contained in the closure of $\pi^{-1}(z)$. By the assumption on the non-degeneracy of \mathcal{C}_x in $\mathbb{P}T_x(X)$, this implies that Z° is a point. It follows that any two general points of X can be connected chain consisting of members of \mathcal{M} of length at most $\dim X$. Let $Y \subset \text{Chow}(X)$ be the

closed subscheme parametrizing connected 1-cycles each component of which is a member of \mathcal{M} . In the notation of Lemma IV.3.4 of [Ko], $u^{(2)} : U \times_Y U \rightarrow X \times X$ is dominant, where $g : U \rightarrow Y$ is the universal family and $u : U \rightarrow X$ is the cycle map. Since U is complete, $u^{(2)}$ is surjective as in the proof of Corollary IV.3.5 of [Ko]. Thus we can apply Proposition 3.13.3 of [Ko] to conclude that $CH_1(X) \otimes \mathbb{Q}$ is generated by members of \mathcal{M} . \square

This proposition can now be used to obtain the Chow generation of one cycles on the moduli space $\mathcal{SU}_C(r, \mathcal{O}_C)$.

Corollary 5.2. *(Hwang) Let $X = \mathcal{SU}_C(r, \mathcal{O}_C)$ for a curve C of genus ≥ 4 . Then $CH_1(X) \otimes \mathbb{Q}$ is generated by Hecke curves. Moreover, there is a surjection*

$$CH_0(C) \otimes \mathbb{Q} \longrightarrow CH_1(\mathcal{SU}_C(r, \mathcal{O}_C)) \otimes \mathbb{Q}.$$

Proof. By Theorem 3 of [Hw2], the union of the tangent vectors to Hecke curves through a general point $y \in X$ is a non-degenerate subvariety $\mathcal{C}_y \subset \mathbb{P}T_y(X)$. Since Hecke curves have degree one with respect to the generator of $\text{Pic}(X)$, they form a closed subscheme \mathcal{M} in $\text{Chow}(X)$. Thus Proposition 5.1 applies. This implies that there is a surjection

$$CH_0(\mathcal{S}) \otimes \mathbb{Q} \rightarrow CH_1(\mathcal{SU}_C(r, \mathcal{O}_C)) \otimes \mathbb{Q}.$$

For any $x \in C$, the moduli space $\mathcal{SU}_C(r, \mathcal{O}(-x))$ is rationally connected and we have the triviality $CH_0(\mathcal{SU}_C(r, \mathcal{O}(-x))) \otimes \mathbb{Q} = \mathbb{Q}$. This gives us an isomorphism

$$CH_0(C) \otimes \mathbb{Q} \simeq CH_0(\mathcal{S}) \otimes \mathbb{Q}.$$

Altogether, we now get a surjection

$$CH_0(C) \otimes \mathbb{Q} \longrightarrow CH_1(\mathcal{SU}_C(r, \mathcal{O}_C)) \otimes \mathbb{Q}.$$

\square

Corollary 5.3. *The Abel-Jacobi map*

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} \longrightarrow \text{Jac}(C) \otimes \mathbb{Q}$$

is an isomorphism, whenever $r \geq 2$ and $g \geq 4$. This extends to an isomorphism

$$CH_1(\mathcal{SU}_C(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} \longrightarrow \text{Jac}(C) \otimes \mathbb{Q}.$$

Proof. By Corollary 5.2, there is a surjection

$$\text{Jac}(C) \otimes \mathbb{Q} \longrightarrow CH_1(\mathcal{SU}_C(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q}.$$

By Corollary 4.2 for $r \geq 2$, the Abel-Jacobi map

$$AJ_1^s : CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} \rightarrow \text{Jac}(C) \otimes \mathbb{Q}$$

is surjective. Furthermore, the localization map

$$CH_1(\mathcal{SU}_C(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q} \rightarrow CH_1(\mathcal{SU}_C^s(r, \mathcal{O}_C))_{\text{hom}} \otimes \mathbb{Q}$$

is also surjective [Fu, p.21]. Putting these facts together, we note that all the maps in the following sequence

$$(17) \quad Jac(C) \otimes \mathbb{Q} \rightarrow CH_1(SU_C(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \rightarrow CH_1(SU_C^s(r, \mathcal{O}_C))_{hom} \otimes \mathbb{Q} \rightarrow Jac(C) \otimes \mathbb{Q}$$

are surjective. Since the composed map $Jac(C) \otimes \mathbb{Q} \rightarrow Jac(C) \otimes \mathbb{Q}$ is surjective, it is an isomorphism. Hence all the maps in (17) are isomorphisms. \square

REFERENCES

- [Ar] Arapura, D. *Letter to the author*, dated 13 April 2009.
- [Ar-Sa] Arapura, D., Sastry, P. *Intermediate Jacobians and Hodge structures of moduli spaces*. Proc. Indian Acad. Sci. Math. Sci. 110 (2000), no. **1**, 1–26.
- [At-Bo] Atiyah, M. F., Bott, R. *The Yang-Mills equations over Riemann surfaces*. Philos. Trans. Roy. Soc. London Ser. A 308 (1983), no. **1505**, 523–615.
- [Ba] Balaji, V. *Cohomology of certain moduli spaces of vector bundles*. Proc. Indian Acad. Sci. Math. Sci. 98 (1988), no. **1**, 1–24.
- [Ba2] Balaji, V. *Intermediate Jacobian of some moduli spaces of vector bundles on curves*. Amer. J. Math. 112 (1990), no. **4**, 611–629.
- [Ba-Se] Balaji, V., Seshadri, C.S. *Cohomology of a moduli space of vector bundles*. The Grothendieck Festschrift, Vol. I, 87–120, Progr. Math., **86**, Birkhuser Boston, Boston, MA, 1990.
- [Ba-Kg-Ne] Balaji, V., King, A. D., Newstead, P. E. *Algebraic cohomology of the moduli space of rank 2 vector bundles on a curve*. Topology 36 (1997), no. **2**, 567–577.
- [Be] Beauville, A. *Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions theta*. (French) [Rank-2 vector bundles over a curve, determinant bundle and theta functions] Bull. Soc. Math. France 116 (1988), no. **4**, 431–448 (1989).
- [BBDe] Beilinson, A. A., Bernstein, J., Deligne, P. *Faisceaux pervers*. (French) [Perverse sheaves] Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astrisque, **100**, Soc. Math. France, Paris, 1982.
- [Bl-Mr] Bloch, S., Murre, J. *On the Chow group of certain types of Fano threefolds*. Compositio Math. 39 (1979), no. **1**, 47–105.
- [Ch-Hw] Choe, I., Hwang, J-M., *Chow group of 1-cycles on the moduli space of vector bundles of rank 2 over a curve*. Math. Z. 253 (2006), no. **2**, 281–293.
- [De] Deligne, P. *Théorie de Hodge. II*. (French) Inst. Hautes Études Sci. Publ. Math. No. **40** (1971), 5–57.
- [Du] Durfee, A. H. *Intersection homology Betti numbers*. Proc. Amer. Math. Soc. 123 (1995), no. **4**, 989–993.
- [Es-Vi] Esnault, H., Viehweg, E. *Deligne-Beilinson cohomology. Beilinson's conjectures on special values of L-functions*, 43–91, Perspect. Math., **4**, Academic Press, Boston, MA, 1988.
- [Fu] Fulton, W. *Intersection theory*, Ergeb. der Math. und ihrer Grenz. 3. Folge, Springer-Verlag, Berlin, 1984.
- [Go-Ma] Goresky, M., MacPherson, R. *Intersection homology theory*. Topology 19 (1980), no. **2**, 135–162.
- [Go-Ma2] Goresky, M., MacPherson, R. *Intersection homology. II*. Invent. Math. 72 (1983), no. **1**, 77–129.
- [Hw-Mo] Hwang, J-M., Mok, N. *Varieties of minimal rational tangents on uniruled projective manifolds. Several complex variables (Berkeley, CA, 1995–1996)*, 351–389, Math. Sci. Res. Inst. Publ., **37**, Cambridge Univ. Press, Cambridge, 1999.
- [Hw] Hwang, J-M. *Tangent vectors to Hecke curves on the moduli space of rank 2 bundles over an algebraic curve*. Duke Math. J. 101 (2000), no. **1**, 179–187.
- [Hw2] Hwang, J-M. *Hecke curves on the moduli space of vector bundles over an algebraic curve*. Proceedings of the symposium, Algebraic Geometry in East Asia (Kyoto, August 3–10, 2001), 155–164, World Scientific, 2003.
- [Ja] Jannsen, U. *Mixed motives and algebraic K-theory*. With appendices by S. Bloch and C. Schoen. Lecture Notes in Mathematics, **1400**. Springer-Verlag, Berlin, 1990. xiv+246 pp.

- [Ki] Kirwan, F. *On the homology of compactifications of moduli spaces of vector bundles over a Riemann surface*. Proc. London Math. Soc. (3) 53 (1986), no. 2, 237–266.
- [Ko] Kollár, J.: *Rational curves on algebraic varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete, 3 Folge, Band 32, Springer Verlag, 1996
- [Na-Ra] Narasimhan, M. S., Ramanan, S. *Deformations of the moduli space of vector bundles over an algebraic curve*. Ann. Math. (2) **101** (1975), 391–417.
- [Na-Ra2] Narasimhan, M. S., Ramanan, S. *Geometry of Hecke cycles. I*. C. P. Ramanujam—a tribute, pp. 291–345, Tata Inst. Fund. Res. Studies in Math., **8**, Springer, Berlin-New York, 1978.
- [Na-Se] Narasimhan, M. S., Seshadri, C. S. *Stable and unitary vector bundles on a compact Riemann surface*, Ann. of Math. (2) **82** 1965 540–567.
- [Ne] Newstead, P. E. *Characteristic classes of stable bundles of rank 2 over an algebraic curve*. Trans. Amer. Math. Soc. **169** (1972), 337–345.
- [Pe-St] Peters, C. A. M.; Steenbrink, J. H. M. *Mixed Hodge structures*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, **52**. Springer-Verlag, Berlin, 2008. xiv+470 pp.
- [Ra] Ramanan, S. *The moduli spaces of vector bundles over an algebraic curve*. Math. Ann. **200** (1973), 69–84.
- [Se] Seshadri, C. S. *Space of unitary vector bundles on a compact Riemann surface*. Ann. of Math. (2) **85** 1967 303–336.
- [Se2] Seshadri, C.S. *Desingularisation of the moduli varieties of vector bundles on curves*. Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977), pp. 155–184, Kinokuniya Book Store, Tokyo, 1978.
- [Se3] Seshadri, C.S. *Fibrés vectoriels sur les courbes algébriques*. (French) [Vector bundles over algebraic curves] Notes written by J-M. Drezet from a course at the École Normale Supérieure, June 1980. Astrisque, **96**. Société Mathématique de France, Paris, 1982. 209 pp.

THE INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI, CHENNAI 600113, INDIA

E-mail address: jniyer@imsc.res.in