

SIGMA THEORY AND TWISTED CONJUGACY-II: HOUGHTON GROUPS AND PURE SYMMETRIC AUTOMORPHISM GROUPS

DACIBERG L. GONÇALVES AND PARAMESWARAN SANKARAN

Abstract Let $\phi : \Gamma \rightarrow \Gamma$ be an automorphism of a group Γ . We say that $x, y \in \Gamma$ are in the same ϕ -twisted conjugacy class and write $x \sim_\phi y$ if there exists an element $\gamma \in \Gamma$ such that $y = \gamma x \phi(\gamma^{-1})$. This is an equivalence relation on Γ called the ϕ -twisted conjugacy. Let $R(\phi)$ denote the number of ϕ -twisted conjugacy classes in Γ . If $R(\phi)$ is infinite for all $\phi \in \text{Aut}(\Gamma)$, we say that Γ has the R_∞ -property.

The purpose of this note is to show that the symmetric group S_∞ , the Houghton groups and the pure symmetric automorphism groups have the R_∞ -property. We show, also, that the Richard Thompson group T has the R_∞ -property. We obtain a general result establishing the R_∞ -property of finite direct product of finitely generated groups.

This is a sequel to an earlier work by Gonçalves and Kochloukova, in which it was shown, using the sigma-theory due to Bieri-Neumann-Strebel, that for most of the groups Γ considered here, $R(\phi) = \infty$ where ϕ varies in a finite index subgroup of the automorphisms of Γ .

1. INTRODUCTION

Let Γ be a group and let $\phi : \Gamma \rightarrow \Gamma$ be an endomorphism. Then ϕ determines an action Φ of Γ on itself where, for $\gamma \in \Gamma$ and $x \in \Gamma$, we have $\Phi_\gamma(x) = \gamma x \phi(\gamma^{-1})$. The orbits of this action are called the ϕ -twisted conjugacy classes. We write $x \sim_\phi y$ if x and y are in the same ϕ -twisted conjugacy class. Note that when ϕ is the identity automorphism, the orbits are the usual conjugacy classes of Γ . We denote by $\mathcal{R}(\phi)$ the set of all ϕ -twisted conjugacy classes and by $R(\phi)$ the cardinality $\#\mathcal{R}(\phi)$ of $\mathcal{R}(\phi)$. We say that Γ has the R_∞ -property if $R(\phi) = \infty$, that is if $\mathcal{R}(\phi)$ is infinite, for every automorphism ϕ of Γ .

The problem of determining which groups have the R_∞ -property—more briefly the R_∞ -problem—has attracted the attention of many

2010 *Mathematics Subject Classification.* 20E45, 20E36.

Keywords and phrases: Twisted conjugacy, Reidemeister number, sigma theory, Houghton groups, infinite symmetric group, pure symmetric automorphism groups.

researchers after it was discovered that all non-elementary Gromov-hyperbolic groups have the R_∞ -property. See [23] and [13]. It is particularly interesting when the group in question is finitely generated or countable. The notion of twisted conjugacy arises naturally in fixed point theory, representation theory, algebraic geometry and number theory. In recent years the R_∞ -problem has emerged as an active research area.

Recall that Houghton introduced a family of groups $H_n, n \geq 2$, defined as follows: Let $M_n := \{1, 2, \dots, n\} \times \mathbb{N}$. The group H_n consists of all bijections $f : M_n \rightarrow M_n$ such that there exists integers t_1, \dots, t_n such that $f(j, s) = (j, s + t_j)$ for all $s \in \mathbb{N}$ sufficiently large and all $j \leq n$. Note that necessarily $\sum_{1 \leq j \leq n} t_j = 0$. Let $Z = \{(t_1, \dots, t_n) \mid \sum_{1 \leq j \leq n} t_j = 0\} \subset \mathbb{Z}^n \cong \mathbb{Z}^{n-1}$. One has a surjective homomorphism $\tau : H_n \rightarrow Z \cong \mathbb{Z}^{n-1}$ sending f to its *translation part* (t_1, \dots, t_n) (with notation as above). It is easily verified that τ is surjective with kernel the group of all *finitary* permutations of M_n . K. S. Brown [6] showed that H_n is finitely presented for $n \geq 3$ and that it is FP_{n-1} but not FP_n . Note that the above definition of H_n makes sense even for $n = 1$ and we have $H_1 \cong S_\infty$. However, we treat the group S_∞ separately and we shall always assume that $n \geq 2$ while considering the family H_n .

Next we recall the group G_n , the group of *pure symmetric automorphisms* of the free group F_n of rank $n \geq 2$. Fix a basis $x_k, 1 \leq k \leq n$, of F_n . Denote by $\alpha_{ij} \in \text{Aut}(F_n), 1 \leq i \neq j \leq n$, the automorphism defined as $x_i \mapsto x_j x_i x_j^{-1}, x_k \rightarrow x_k, 1 \leq k \leq n, k \neq i$. The group G_n is the subgroup of $\text{Aut}(F_n)$ generated by $\alpha_{ij}, 1 \leq i \neq j \leq n$. McCool showed that G_n is finitely presented where the generating relations are:

- (i) $[\alpha_{ij}, \alpha_{kl}] = 1$, whenever i, j, k, l are all different;
- (ii) $[\alpha_{ik}, \alpha_{jk}] = 1$ and $[\alpha_{ij}\alpha_{kj}, \alpha_{ik}] = 1$ whenever i, j, k are all different.

It was shown by Gonçalves and Kochloukova [15] that $R(\phi) = \infty$ for all ϕ in a finite index subgroup of the group of all automorphisms of Γ where $\Gamma = H_n, G_n$. Our main result is the following theorem. We give two proofs for the case of Houghton groups, neither of which use Σ -theory. However we still need to use the results of [15] in the case of G_n .

Theorem 1.1. *The following groups have the R_∞ -property:*

- (i) *The group S_∞ of finitary permutations of \mathbb{N} ,*
- (ii) *the Houghton groups $H_n, n \geq 2$, and,*
- (iii) *the group $G_n, n \geq 2$, of pure symmetric automorphisms of a free group of rank n .*

Recall that Richard Thompson constructed three groups finitely presented infinite groups $F \subset T \subset V$ around 1965 and showed that T and V are simple. The groups F, T , and V arise as certain homeomorphism groups of the reals, the circle, and the Cantor set respectively. Since then these constructions have been generalized by G. Higman [21]. See also K. S. Brown [6], R. Bieri and R. Strebel [3], and M. Stein [29]. For an introduction to the Thompson groups F, T, V see [10].

Theorem 1.2. *The Richard Thompson group T has the R_∞ -property.*

As the group T is simple, Σ -theory yields no information about the R_∞ -property. The above theorem was first proved by Burillo, Matucci, and Ventura [8]. Shortly thereafter, Gonçalves and Sankaran [16] also independently obtained the same result.

In section §2 we make some preliminary observations concerning the R_∞ -property which will be needed for our purposes. Theorem 1.1 will be established in §3. The R_∞ -property of the group T will be proved in §4. In §5 we consider the R_∞ -property of finite direct product of groups and obtain a strengthening of a result of Gonçalves and Kochloukova [15].

This is a sequel to the paper [15] by Gonçalves and Kochloukova. We reassure the reader that this paper can be read independently of it. Although results from [15] are used, we develop our own proof techniques to go forward.

Note. Just after this paper was submitted, J. H. Jo, J. B. Lee, and S. R. Lee [22] have announced almost simultaneously a proof of the R_∞ property for the Houghton groups.

If $f : X \rightarrow Y$ is a map of sets, we shall always write the argument to the right of f ; thus $f(x)$ denotes the image of $x \in X$ under f .

2. PRELIMINARIES

We begin by recalling some general results concerning twisted conjugacy classes of an automorphism of a group and that of its restriction to a normal subgroup. We obtain a criterion for a periodic automorphism to have infinitely many twisted conjugacy classes. We shall also briefly recall the notion of the Bieri-Neumann-Strebel invariant and give its known description in the case of Houghton groups and the pure symmetric automorphism groups.

2.1. Addition formula. The following addition formula may be found in [17, Lemma 2.1]. This is a special case of a more general formula proved in [18, §2]. For any element $g \in G$, we shall denote by ι_g the inner automorphism $x \mapsto gxg^{-1}$ of G . When N is a normal subgroup

of G , we shall abuse notation and denote by the same symbol ι_g the automorphism of N got by restriction of ι_g to N .

Lemma 2.1. *Suppose that we have a commutative diagram of homomorphisms of groups where the vertical arrows are isomorphisms and horizontal rows are short exact sequence:*

$$\begin{array}{ccccccccc} 1 & \rightarrow & N & \xrightarrow{i} & G & \xrightarrow{p} & G/N & \rightarrow & 1 \\ & & \downarrow \theta' & & \downarrow \theta & & \downarrow \bar{\theta} & & \\ 1 & \rightarrow & N & \xrightarrow{i} & G & \xrightarrow{p} & G/N & \rightarrow & 1. \end{array}$$

Then:

- (i) One has an exact sequence of (pointed) sets $\mathcal{R}(\theta') \xrightarrow{i_*} \mathcal{R}(\theta) \xrightarrow{p_*} \mathcal{R}(\bar{\theta}) \rightarrow \{0\}$. That is, p_* is surjective and $\text{Im}(i_*) = p_*^{-1}(\{N\})$.
- (ii) (Addition formula): Suppose that $R(\bar{\theta}) < \infty$ and that $\text{Fix}(\iota_{\alpha N} \circ \bar{\theta}) = \{N\}$ for all $\alpha \in G$. Then $R(\theta) < \infty$ if and only if $R(\iota_{\alpha} \theta') < \infty$ for all $\alpha \in G$. Moreover, the following addition formula holds if $R(\theta) < \infty$: $R(\theta) = \sum_{[\alpha N] \in \mathcal{R}(\bar{\theta})} R(\iota_{\alpha} \theta')$. \square

We omit the proof. Part (i) is trivial. As mentioned above, the addition formula is also a known result. In any case it can be proved in a straightforward manner. It can also be proved easily using the fixed point version the following six term exact sequence of sets due to P. R. Heath [20, (2), p.4] (cf. [19, Theorem 1.8]):

$$1 \rightarrow \text{Fix}(\iota_{\alpha} \theta') \rightarrow \text{Fix}(\iota_{\alpha} \theta) \rightarrow \text{Fix}(\iota_{\bar{\alpha}} \bar{\theta}) \rightarrow \mathcal{R}(\iota_{\alpha} \theta') \rightarrow \mathcal{R}(\iota_{\alpha} \theta) \rightarrow \mathcal{R}(\iota_{\bar{\alpha}} \bar{\theta}) \rightarrow 1$$

where $\bar{\alpha}$ denotes $\alpha N \in G/N$.

Remark 2.2. Note that if $G/N \cong \mathbb{Z}^n, n < \infty$, and if 1 is not an eigenvalue of the matrix of $\bar{\theta}$ with respect to a basis of G/N , then for any $\alpha \in G, \text{Fix}(\iota_{\alpha N} \circ \bar{\theta}) = \text{Fix}(\bar{\theta})$ consists only of the trivial element. So the lemma implies that, if $R(\theta') = \infty$, then $R(\theta) = \infty$.

2.2. Periodic outer automorphisms. Let Γ be a group with infinitely many conjugacy classes. Then, for any automorphism $\phi : \Gamma \rightarrow \Gamma$, and any $g \in G, R(\phi) = R(\iota_g \circ \phi)$ where ι_g denotes the inner automorphism $x \mapsto gxg^{-1}$. Indeed it is readily seen that the ϕ -twisted conjugacy classes are the same as the left translation by g of the $\iota_g \circ \phi$ -twisted conjugacy classes. Thus Γ has the R_{∞} -property if and only if $R(\phi) = \infty$ for a set of coset representatives of $\text{Out}(\Gamma) = \text{Aut}(\Gamma)/\text{Inn}(\Gamma)$. We have the following lemma. Compare [16, Lemma 3.4].

Lemma 2.3. *Let $\theta \in \text{Aut}(\Gamma)$ and let $n \geq 1$. Suppose that $\{x^n \mid x \in \text{Fix}(\theta)\}$ is not contained in the union of finitely many θ^n -twisted conjugacy classes of Γ . Then $R(\theta) = \infty$.*

Proof. Let $x \sim_\theta y$ in Γ where $x, y \in \text{Fix}(\theta)$. Thus there exists an $z \in \Gamma$ such that $y = z^{-1}x\theta(z)$. Applying θ^i both sides, we obtain $y = \theta^i(z^{-1})x\theta^{i+1}(z)$ as $x, y \in \text{Fix}(\theta)$. Write $\phi := \theta^n$. Multiplying these equations successively for $0 \leq i < n$, we obtain

$$y^n = \prod_{0 \leq i < n} \theta^i(z^{-1})x\theta^{i+1}(z) = z^{-1}x^n\theta^n(z) = z^{-1}x^n\phi(z).$$

That is, $y^n \sim_\phi x^n$. Our hypothesis says that there are infinitely many elements $x_k \in \text{Fix}(\theta)$, $k \geq 1$, such that the x_k^n are in pairwise distinct ϕ -twisted conjugacy classes of Γ . Hence we conclude that $R(\theta) = \infty$. \square

Remark 2.4. When $\theta^n = \iota_\gamma$ is an inner automorphism, we see from the above lemma that $R(\theta) = \infty$ if $\{x^n\gamma \mid x \in \text{Fix}(\theta)\}$ is not contained in a finite union of conjugacy classes of Γ . If $\theta^n = id$, then $R(\theta) = \infty$ if $\text{Fix}(\theta)$ contains elements of order k for arbitrarily large values of $k \in \mathbb{N}$.

2.3. Σ -theory of H_n and G_n . Bieri, Neumann, and Strebel [2] introduced, for any finitely generated group Γ , an invariant $\Sigma(\Gamma)$ which is a certain open subset—possibly empty—of the character sphere $S(\Gamma) := \text{Hom}(\Gamma, \mathbb{R}) \setminus \{0\}/\mathbb{R}_{>0}$ where the action of the multiplicative group of positive reals is via scalar multiplication. The automorphism group $\text{Aut}(\Gamma)$ acts on $S(\Gamma)$ where $\phi^* : S(\Gamma) \rightarrow S(\Gamma)$ is defined as $[\chi] \mapsto [\chi \circ \phi]$, $[\chi] \in S(\Gamma)$, for $\phi \in \text{Aut}(\Gamma)$. This action preserves the subspace $\Sigma(\Gamma)$ and hence also its complement $\Sigma^c(\Gamma)$. If the image of the anti homomorphism $\eta : \text{Aut}(\Gamma) \rightarrow \text{Homeo}(\Sigma^c(\Gamma))$ defined as $\phi \mapsto \phi^*$ is a finite group, then $K = \ker(\eta)$ is a finite index subgroup of $\text{Aut}(\Gamma)$ which fixes every character class in $\Sigma^c(\Gamma)$. This happens, for example, if $\Sigma^c(\Gamma)$ is a non-empty finite set. If $\Sigma^c(\Gamma)$ contains a *discrete* character class $[\chi]$, that is, a class represented by an character χ whose image $\chi(\Gamma) \subset \mathbb{R}$ is infinite cyclic, then it was observed by Gonçalves and Kochloukova [15] that the character χ itself is fixed by the action of K on $\text{Hom}(\Gamma, \mathbb{R})$. That is, $\chi \circ \phi = \chi$ for all $\phi \in K \subset \text{Aut}(\Gamma)$. This easily implies that $R(\phi) = \infty$ by Lemma 2.1(i), taking $G = \Gamma$, $N = \ker \chi$, $\theta = \phi$ in the notation of that lemma, so that $\bar{\theta} = id$.

When $\Gamma = G_n$, $n \geq 3$, the group of pure symmetric automorphisms of F_n , L. Orlandi-Korner [27] has determined $\Sigma^c(\Gamma)$. When $\Gamma = H_n$, the Houghton group, Brown [7] computed the set $\Sigma^c(\Gamma)$. Using these results, Gonçalves and Kochloukova, showed that if Γ is any one of the groups H_n , $n \geq 2$, G_m , $m \geq 3$, the image of $\eta : \text{Aut}(\Gamma) \rightarrow \text{Homeo}(\Sigma^c(\Gamma))$ is finite.

In the case of the Houghton group H_n , $n \geq 2$, it turns out that $\Sigma^c(H_n)$ is a finite set of discrete character classes $[\chi_j]$, $1 \leq j \leq n$. Explicitly, $\chi_j : H_n \rightarrow \mathbb{Z}$ may be taken to be $-\pi_i \circ \tau$ where $\tau : H_n \rightarrow Z$

is the translation part (see §1) and $\pi_i : Z \rightarrow \mathbb{Z}$ is the restriction to $Z \subset \mathbb{Z}^n$ of the i -th projection (see [7]). (Recall from §1 that $Z = \{(t_1, \dots, t_n) \in \mathbb{Z}^n \mid \sum_{1 \leq j \leq n} t_j = 0\}$.) Thus $\text{Homeo}(\Sigma^c(H_n)) \cong S_n$ is finite and so is the image of $\eta : \text{Aut}(H_n) \rightarrow \text{Homeo}(\Sigma^c(H_n))$. As already remarked $R(\phi) = \infty$ for all $\phi \in \ker(\eta)$. The lemma below will not be used in the sequel but included here for illustrative purposes.

Lemma 2.5. *Suppose that $\eta(\phi) : \Sigma^c(H_n) \rightarrow \Sigma^c(H_n)$ is not an n -cycle. Then $R(\phi) = \infty$.*

Proof. Since $\eta(\phi)$ is not an n -cycle, the orbit of $[\chi_1]$ under $\eta(\phi)$ consists of at most $n - 1$ elements. Since χ_1 is discrete, the orbit of $\chi_1 \in \text{Hom}(H_n, \mathbb{R})$ consists of discrete elements. In fact the orbit of χ_1 is a subset of $\{\chi_j \mid 1 \leq j \leq n\}$. Now the orbit sum $\lambda := \sum_{1 \leq j \leq k} \chi_1 \phi^j$ is a *non-zero* character since any $n - 1$ elements of $\chi_j, 1 \leq j \leq n$ form a *basis* of $\text{Hom}(H_n, \mathbb{R})$. It follows that, since $\phi^*(\lambda) = \lambda$, $R(\phi) = \infty$. \square

If $\phi^* : \Sigma^c(H_n) \rightarrow \Sigma^c(H_n)$ is an n -cycle, then the orbit sum is zero and the above argument fails. In fact, it is easily seen that every possible permutation of $\Sigma^c(H_n)$ may be realized as $\eta(\phi)$ for some $\phi \in \text{Aut}(H_n)$, that is, $\eta : \text{Aut}(H_n) \rightarrow \text{Homeo}(\Sigma^c(H_n)) \cong S_n$ is surjective.

3. PROOF OF THEOREM 1.1

Let X be an infinite set. We will only be concerned with the case when X is countably infinite. We shall denote by $S_\infty(X)$ the group of all finitary permutations of X , that is those permutations which fixes all but finitely many elements of X . The group of *all* permutations of X will be denoted by $S(X)$. We shall denote $S(X)$ (resp. $S_\infty(X)$) simply by S_ω (resp. S_∞) when X is clear from the context. If $x = (x_k)_{k \in \mathbb{Z}}$ is a doubly infinite sequence in X of pairwise distinct elements, we regard it as an element of $S(X)$ where $x(x_k) = x_{k+1}$ and $x(a) = a$ if $a \neq x_k \forall k \in \mathbb{Z}$. Two such sequences $x = (x_k)$ and $y = (y_k)$ define the same permutation if and only if y is a shift of x , that is, there exists an n such that $x_k = y_{k+n}$ for all $k \in \mathbb{Z}$. Thus, the sequence $x = (x_k)_{k \in \mathbb{Z}}$ is just the infinite cycle $x \in S(X)$. Any $f \in S(X)$ is uniquely expressible as a product of disjoint cycles. Such an expression of f is its *cycle decomposition*. The *cycle type* of an $f \in S(X)$ is the function $c(f) : \mathbb{N} \cup \{\infty\} \rightarrow \mathbb{Z}_{\geq 0} \cup \{\infty\}$ where $c(f)(\alpha)$ is the number of α -cycles in the cycle decomposition of f if that number is finite, otherwise it is ∞ for $\alpha \in \mathbb{N} \cup \{\infty\}$. As in the case $S_\infty(X)$, if f and g have the same cycle type, then they are conjugate in $S(X)$. We need a criterion for f and g to be conjugate by an element of $S_\infty(X)$.

Lemma 3.1. *Let $x = (x_k)_{k \in \mathbb{Z}}, y = (y_k)_{k \in \mathbb{Z}} \in S_\omega(X)$ be two disjoint infinite cycles and let $(a, b) \in S_\infty$.*

(i) If $a = x_0, b = x_k, k > 0$, then $(a, b)x = uv$, a product of disjoint cycles $u = (u_j)_{j \in \mathbb{Z}} \in S_\omega, v \in S_\infty$, defined as

$$u_j = \begin{cases} x_j, & j < 0, \\ x_{j+k}, & j \geq 0, \end{cases}$$

and $v = (x_0, \dots, x_{k-1}) \in S_\infty$.

(ii) If $a = x_0, b = y_0$, then $(a, b)xy = uv$, where $u = (u_j)_{j \in \mathbb{Z}}, v = (v_j)_{j \in \mathbb{Z}}$ are disjoint infinite cycles defined as

$$u_j = \begin{cases} x_j, & j < 0, \\ y_j, & j \geq 0, \end{cases}$$

and

$$v_j = \begin{cases} y_j, & j < 0, \\ x_j, & j \geq 0. \end{cases}$$

□

If $k \in \mathbb{N}$, we denote by $\mathbb{N}_{>k}$ the set of all integers greater than k . Note that $S_\infty = \cup_{k \geq 2} S_k$ where S_k is the subgroup consisting of permutations of \mathbb{N} which fixes all $n > k$. In particular, the group S_∞ is generated by transpositions $(i, i+1), i \geq 1$. The alternating group A_∞ equals the commutator subgroup $[S_\infty, S_\infty]$, has index 2 in S_∞ and is simple. The conjugacy class of any element of S_∞ is determined by its cycle type, as in the case of finite symmetric groups. The group S_∞ is a normal subgroup of $S_\omega = S(\mathbb{N})$. In particular, any bijection $f : \mathbb{N} \rightarrow \mathbb{N}$ defines an automorphism $\iota_f \in \text{Aut}(S_\infty)$ by restricting the inner automorphism determined by $f \in S_\omega$. Moreover ι_f is the identity automorphism only if f equals the identity map. The following result is well-known. See [28, §11.4].

Theorem 3.2. *The homomorphism $\iota : S_\omega \rightarrow \text{Aut}(S_\infty)$ is an isomorphism of groups.*

The following corollary is a special case of a more general result established in [12, Theorem 8.2A]. We include a proof, which is simpler in our special case.

Corollary 3.3. *Suppose that S_∞ is a characteristic subgroup of a group H contained in S_ω . Then the automorphism group of H is isomorphic to the normalizer of $N(H)$ of H in S_ω . In particular, every automorphism of H is the restriction to H of a unique inner automorphism of S_ω .*

Proof. We shall denote by the same symbol ι_f to denote the conjugation by $f \in S_\omega$ or its restriction to any subgroup normalized by f .

It is evident that $\iota : N(H) \rightarrow \text{Aut}(H)$ defined as $f \mapsto \iota_f$ defines an homomorphism. (Here $\iota_f(h) = fhf^{-1} \forall h \in H$.) This is a monomorphism since ι_f is non-trivial on $S_\infty \subset H$ if f is not the identity.

Let $\phi : H \rightarrow H$ be any automorphism and let $f \in S_\omega$ be the element such that $\phi|_{S_\infty} = \iota_f$. We claim that $\phi = \iota_f$. Suppose that $u := \phi(h), \iota_f(h) = fhf^{-1} =: v$ for some $h \in H$. We must show that $u(i) = v(i)$ for all $i \in \mathbb{N}$. It suffices to show that $\{u(i), u(j)\} = \{v(i), v(j)\}$ for all $i, j \in \mathbb{N}, i \neq j$. Let $i, j \in \mathbb{N}, i \neq j$. Now consider the transposition $(a, b) \in S_\infty$ such that $\iota_f(a, b) = \phi(a, b) = (i, j)$. We have $\phi(h(a, b)h^{-1}) = \phi(h)\phi(a, b)\phi(h^{-1}) = u(i, j)u^{-1} = (u(i), u(j))$, while $\iota_f(h(a, b)h^{-1}) = \iota_f(h)\iota_f(a, b)\iota_f(h^{-1}) = v(i, j)v^{-1} = (v(i), v(j))$. Therefore $(u(i), u(j)) = (v(i), v(j)) \in S_\infty$ since ι_f and ϕ agree on S_∞ . This implies that $\{u(i), u(j)\} = \{v(i), v(j)\}$, completing the proof. \square

3.1. S_∞ has the R_∞ -property. Let $\theta \in \text{Aut}(S_\infty)$. In view of Theorem 3.2 $\theta = \iota_f$ for some $f \in S_\omega$. Let $x, y \in S_\infty$ and suppose that $y = zx\theta(z^{-1}) = zxfz^{-1}f^{-1}$ for some $z \in S_\infty$. Then $yf = z(xf)z^{-1}$ in S_ω for some $z \in S_\infty$. For any cycle (finite or infinite) $u = (u_j)$, we have that zuz^{-1} is the cycle $(z(u_j))$. Any $z \in S_\infty$ moves only finitely many elements of \mathbb{N} . Hence when u is an infinite cycle $z(u_j) = u_j$ for all but finitely many $j \in \mathbb{Z}$. For an arbitrary element u expressed as a product of pairwise disjoint cycles, $u(\alpha) = (u(\alpha)_j)$, the element zuz^{-1} being a product of $zu(\alpha)z^{-1}$, we see that $zu(\alpha)z^{-1} = u(\alpha)$ for all but a finitely many α , and, moreover, if $u(\alpha) = (u(\alpha)_j)_{j \in \mathbb{Z}}$ is an infinite cycle, then $z(u(\alpha)_j) = u(\alpha)_j$ for all but finitely many $j \in \mathbb{Z}$.¹

Lemma 3.4. *Suppose that $f \in S_\omega$ has an infinite cycle u , then there exist infinitely many transpositions $\tau_k \in S_\infty$ such that $\tau_j f \neq z\tau_k f z^{-1}$ for any $z \in S_\infty$.*

Proof. Fix an infinite cycle $u = (u_\alpha)_{\alpha \in \mathbb{Z}}$ that occurs in the cycle decomposition of f . Let $\tau_\alpha = (u_0, u_\alpha), \alpha \geq 1$. Then we claim that $\tau_\alpha f$ and $\tau_\beta f$ are not conjugates if $\alpha \neq \beta$. To see this, we apply Lemma 3.1 to compute $\tau_\alpha u, \alpha \geq 1$. Note that the cycles that occur in $\tau_\alpha u$ also occur in the cycle decomposition of $\tau_\alpha f$. This is true in particular of the infinite cycle, denoted $v(\alpha)$, that occurs in $\tau_\alpha u$.

Now $v(\alpha)_p = v(\beta)_p = u_p$ for all $p < 0$ and $\alpha, \beta \geq 1$, and, when $\alpha \neq \beta$, $u_{p+\alpha} = v(\alpha)_p \neq v(\beta)_p = u_{p+\beta}, p \geq 0$. This implies that the $zv(\beta)z^{-1}$ cannot occur in $\tau_\alpha f$ for any $z \in S_\infty$ if $\alpha \neq \beta$ in its cycle decomposition

¹There is a mild abuse of notation here; $u(\alpha)$ is not to be confused with the value of u at α . We will use Greek letters as labels in such situations.

by the assertion made in the paragraph above the statement of the lemma. Hence $\tau_\alpha f \neq z\tau_\beta f z^{-1}$ for any $z \in S_\infty$. \square

We are now ready to prove the first assertion of Theorem 1.1.

Theorem 3.5. *The group S_∞ has the R_∞ -property.*

Proof. Let $\theta = \iota_f \in \text{Aut}(S_\infty)$ where $f \in S_\omega$. We need to show that there exists pairwise distinct elements $\tau_j \in S_\infty, j \in \mathbb{N}$ such that $\tau_j f \neq z\tau_k f z^{-1}$ for any $z \in S_\infty$ if $j \neq k$. Since S_∞ has infinitely many conjugacy classes, the assertion holds for $f \in S_\infty$, we need only consider the case $f \notin S_\infty$. Thus, in the cycle decomposition of f , either (i) there exist an infinite cycle, or, (ii) all the cycles are finite and there are infinitely many of them.

Case (i). In this case the assertion has already been established in Lemma 3.4.

Case(ii). Suppose that $f = \prod_{\alpha \in \mathbb{N}} u(\alpha)$ where the $u(\alpha)$ are all finite cycles having length $\ell(\alpha)$ at least 2 for every $\alpha \in \mathbb{N}$. Let $J := \{\alpha \in \mathbb{N} \mid \ell(\alpha) \geq 3\}$. We break up the proof into two subcases depending on whether J is infinite or not.

Subcase (a): J is infinite. Let $J_k \subset J$ be the set consisting of the first k elements of J (with respect to the usual ordering on $J \subset \mathbb{N}$). Write $u(\alpha) = (u(\alpha)_1, \dots, u(\alpha)_{\ell(\alpha)})$ and set $U(\alpha) := \{u(\alpha)_i \mid 1 \leq i \leq \ell(\alpha)\}, \alpha \in \mathbb{N}$. Consider the collection of pairwise disjoint transpositions $\lambda_\alpha = (u(\alpha)_1, u(\alpha)_2), \alpha \in J$, and let $\tau_k = \prod_{\alpha \in J_k} \lambda_\alpha$. Note that $\lambda_\alpha u(\alpha) = (u(\alpha)_1) \cdot (u(\alpha)_2, \dots, u(\alpha)_{\ell(\alpha)}) = (u(\alpha)_2, \dots, u(\alpha)_{\ell(\alpha)})$ fixes only $u(\alpha)_1$ in the set $U(\alpha)$ as $\ell(\alpha) \geq 3$. Then $\tau_k \cdot \prod_{\alpha \in J_k} u(\alpha)$ fixes only $u(\alpha)_1 \in \mathbb{N}, \alpha \in J_k$, in the set $\cup_{\alpha \in J_k} U(\alpha)$. Let $F_0 = \text{Fix}(f)$. Then $\text{Fix}(\tau_k f) = F_0 \cup \{u(\alpha)_1 \mid \alpha \in J_k\} =: F_k$.

Suppose that $\tau_j f = z\tau_k f z^{-1}$ with $z \in S_\infty$ with $j \neq k$. Then z defines a bijection $\zeta : F_j \rightarrow F_k$ between the fixed sets of $\tau_j f$ and $\tau_k f$. Clearly this is a contradiction if $\text{Fix}(f) = F_0$ is finite. Assume that $F_0 \subset \mathbb{N}$ is infinite. Since $z \in S_\infty$, it fixes all but finitely many elements of F_0 . Let $L := \{m \in F_0 \mid z(m) \neq m\}$. Note that ζ restricts to the identity on $F_0 \setminus L$. Therefore ζ restricts to a bijection between $L \cup \{u(\beta)_1 \mid \beta \in J_j\}$ and $L \cup \{u(\beta)_1 \mid \beta \in J_k\}$. Since $j \neq k$, L is finite and $L \subset F_0$ is disjoint from $\{u(\beta)_1 \mid \beta \in J_n\}, n = j, k$, this is a contradiction.

Subcase (b): The set J is finite; we set $K = \mathbb{N} \setminus J$ and define $K_j, j \in \mathbb{N}$, to be the set of first α elements of K . Again we set $\lambda_\alpha = (u(\alpha)_1, u(\alpha)_2) = u(\alpha), \alpha \in K$. Now, if $\alpha \in K$, we have $\lambda_\alpha u(\alpha) = id$, that is, $\lambda_\alpha u(j)$ fixes both points of $U(\alpha)$. We set $\tau_j := \prod_{\alpha \in K_j} \lambda_\alpha, F_j := \text{Fix}(\tau_j f) = F_0 \cup_{\alpha \in K_j} U(\alpha)$. Arguing exactly as above we see that for any $z \in S_\infty, \tau_j f = z\tau_k f z^{-1}$ implies $j = k$, completing the proof. \square

3.2. Houghton groups. As in the introduction, $H_n, n \geq 2$, denotes the Houghton group. We first describe the group of outer automorphisms of H_n . Recall from §1 that one has an exact sequence

$$1 \rightarrow S_\infty(M_n) \hookrightarrow H_n \xrightarrow{\tau} Z \rightarrow 1$$

where $\tau : H_n \rightarrow Z$ send $f \in H_n$ to the translation part $(t_1, \dots, t_n) \in Z$ of f . The group $S_\infty(M_n)$ is the commutator subgroup of H_n if $n \geq 3$. When $n = 2$, the commutator subgroup is the alternating group $A_\infty(M_2)$ which has index 2 in $S_\infty(M_2)$. In any case, $S_\infty = S_\infty(M)$ is characteristic in H_n as H_n/S_∞ is the maximal *torsion-free* abelian quotient of H_n .

Lemma 3.6. *Let $\phi : H_n \rightarrow H_n, n \geq 2$, be an automorphism. Then ϕ is inner if and only if $\bar{\phi} : Z \rightarrow Z$ is the identity automorphism.*

Proof. It is trivial to see that any inner automorphism of H_n induces the identity automorphism of Z . For the converse, suppose that $\phi : H_n \rightarrow H_n$ induces the identity automorphism of Z .

Let $f \in S(M_n)$ be such that $\iota_f(H_n) = H_n$.

Consider the element $h_p : M_n \rightarrow M_n, 1 \leq p < n$, in H_n defined follows: ²

$$h_p(i, k) = \begin{cases} (p, k+1), & \text{if } i = p, k \geq 1 \\ (n, k-1), & \text{if } i = n, k > 1, \\ (p, 1), & \text{if } i = n, k = 1 \\ (i, k), & \text{if } i \neq p, n. \end{cases}$$

Thus h_p permutes $\{p, n\} \times \mathbb{N}$ in a single cycle:

$$h_p = (\dots, (n, 2), (n, 1), (p, 1), (p, 2), \dots, (p, k), \dots)$$

and so $fh_p f^{-1}$ is the cycle

$$fh_p f^{-1} = (\dots, f(n, 2), f(n, 1), f(p, 1), f(p, 2), \dots, f(p, k), \dots) \in H_n.$$

The only infinite cycles in H_n are those whose terms, except for a finite part of the cycle, are *consecutive* numbers along two rays, say $\{i_n\} \times \mathbb{N}$ and $\{i_p\} \times \mathbb{N}$, in the negative and positive directions respectively of the cycle $fh_p f^{-1}$. Therefore we have $\tau(fh_p f^{-1}) = e_{i_p} - e_{i_n}$. Moreover, there exist integers t_n, t_p such that $f(n, k) = (i_n, k + t_n), f(p, k) = (i_p, k + t_p)$ for sufficiently large k . Clearly i_n and t_n are independent of p . Since f is a bijection, the association $p \mapsto i_p$ is a permutation $\pi_f \in S_n$, and $\sum_{1 \leq q \leq n} t_q = 0$. Note that $\pi_f = id$ if and only if $f \in H_n$.

Since S_∞ is characteristic in H_n , by Corollary 3.3, $\phi = \iota_g$ for a unique $g \in S(M_n)$. We claim that $g \in H_n$. Since $\tau(ghg^{-1}) = \tau(\phi(h)) =$

²The element $(p, k) \in M_n$ should not be confused with the transposition in $S(\mathbb{N})$.

$\tau(h) \forall h \in H_n$, we have $\pi_g(q) = q$ for all $q \leq n$ and so we have $g \in H_n$. \square

The group S_n acts on the set $M_n = \{1, \dots, n\} \times \mathbb{N}$ in the obvious manner, by acting via the identity on \mathbb{N} . This defines an action ψ of S_n on the group $S(M_n)$ defined as $f \mapsto \sigma \circ f \circ \sigma^{-1}$ which preserves the subgroup H_n . Thus we obtain a homomorphism $\psi : S_n \rightarrow \text{Aut}(H_n)$. It is readily seen that $\tau(\psi_\sigma(h)) = \sigma(\tau(h)) \forall h \in H_n$, where σ acts on $Z \subset \mathbb{Z}^n$ by permuting the standard basis elements e_1, \dots, e_n . In particular ψ is a monomorphism. Let $\bar{\psi} : S_n \rightarrow \text{Out}(H_n)$ be the composition of ψ with the projection $\text{Aut}(H_n) \rightarrow \text{Out}(H_n)$.

Proposition 3.7. The homomorphism $\bar{\psi} : S_n \rightarrow \text{Out}(H_n)$ is an isomorphism and so $\text{Aut}(H_n) = \text{Inn}(H_n) \rtimes S_n \cong H_n \rtimes S_n$.

Proof. Lemma 3.6 shows that $\bar{\psi}$ is a monomorphism. We shall show that it is surjective.

Let $\phi \in \text{Aut}(H_n)$. Write $\phi = \iota_f$ for a (unique) $f \in S(M_n)$. With notations as in the proof of Lemma 3.6, let $\pi := \pi_f \in S_n$.

Consider the automorphism $\psi_\pi^{-1}\phi =: \theta$. We have $\tau(\theta(h_p)) = \pi^{-1}(\tau(\phi(h_p))) = \pi^{-1}(\tau(fh_p f^{-1})) = \pi^{-1}(e_{\pi(p)} - e_{\pi(n)}) = e_p - e_n = \tau(h_p), 1 \leq p < n$. Since the group Z is generated by $\tau(h_p), 1 \leq p < n$, it follows by Lemma 3.6 that θ is inner. Hence $\bar{\psi}(\pi) = \phi \pmod{\text{Inn}(H_n)}$.

Finally note that $\text{Inn}(H_n) \cong H_n$ since the centre of H_n is trivial. \square

The above description of $\text{Aut}(H_n)$ had been obtained by Burillo, Cleary, Martino, and Röver [9, Theorem 2.2] and also by Cox [11, §2.2]. All the proofs make essential use of Theorem 3.2 and Corollary 3.3. The proof given by Burillo et al. and our proof seem to be based on the same idea, although conceived of independently.

Theorem 3.8. *The Houghton group H_n has the R_∞ -property for any $n \geq 2$.*

We shall give two proofs. The first one uses the structure of the automorphism group of H_n and is more direct. The second one uses the result of Theorem 3.5 and the addition formula (Lemma 2.1).

First proof. First observe that there are infinitely many conjugacy classes in H_n since two elements in $S_\infty = S_\infty(M_n) \subset H_n$ are conjugates in H_n only if they have the same cycle type. It follows that $R(\phi) = \infty$ for any inner automorphism ϕ of H_n . Therefore, to show that $R(\phi) = \infty$ for an arbitrary $\phi \in \text{Aut}(H_n)$, it suffices to show that $R(\phi) = \infty$ for all ϕ in a set of coset representatives of elements of $\text{Out}(H_n)$. Thus we need only show that $R(\psi_\sigma) = \infty$ for any $\sigma \in S_n$, where $\psi : S_n \rightarrow$

$\text{Aut}(H_n)$ is as defined in the paragraph above Proposition 3.7. We shall use Lemma 2.3 and Remark 2.4 to achieve this.

For $k \geq 1$, consider the element ξ_k defined as the product of k -cycles $((i, 1), \dots, (i, k)) \in H_n$, $1 \leq i \leq n$. Explicitly,

$$\xi_k(i, j) = \begin{cases} (i, j+1) & \text{if } 1 \leq j < k, \\ (i, 1) & \text{if } j = k, \\ (i, j) & \text{if } j > k, \end{cases}$$

for all $i \leq n$. Then ξ_k is fixed by ψ_σ for every $\sigma \in S_n$. Thus, $\{\xi_k^n \mid k \geq 1\}$ contains elements of arbitrarily large orders and so by Remark 2.4 it follows that $R(\psi_\sigma) = \infty$ for all $\sigma \in S_n$, completing the proof.

Second proof. Consider the exact sequence $1 \rightarrow S_\infty(M_n) \rightarrow H_n \rightarrow Z \rightarrow 0$. As remarked already, $S_\infty(M_n)$ is characteristic in H_n and we have $Z \cong \mathbb{Z}^{n-1}$. Thus any automorphism θ of H_n restricts to an automorphism θ' of $S_\infty(M_n)$ and induces an automorphism $\bar{\theta}$ of Z . If $R(\bar{\theta}) = \infty$ then, by Lemma 2.1 (i), we have $R(\theta) = R(\bar{\theta}) = \infty$. Now suppose that $R(\bar{\theta}) < \infty$. Then $\text{Fix}(\bar{\theta}) = 0$. Since Z is abelian and since $R(\theta') = \infty$ by Theorem 3.5, the addition formula (Lemma 2.1(ii)) yields $R(\theta) = R(\theta') = \infty$, completing the proof. \square

3.3. The group of pure symmetric automorphisms. Recall that $G_n \subset \text{Aut}(F_n)$, $n \geq 2$, denotes the group of pure symmetric automorphisms of the free group F_n of rank n . A presentation for G_n , obtained by McCool, was recalled in §1. It is immediate from this presentation that the abelianization $G_n^{\text{ab}} = G_n/[G_n, G_n]$ is isomorphic to \mathbb{Z}^{n^2-n} with basis the images $\bar{\alpha}_{ij}$, $1 \leq i \neq j \leq n$. We denote by $\{\chi_{ij} \mid 1 \leq i \neq j \leq n\}$ the basis of $\text{Hom}(G_n^{\text{ab}}, \mathbb{Z})$ dual to the basis $\{\bar{\alpha}_{ij} \mid 1 \leq i \neq j \leq n\}$. We shall denote by the same symbol χ_{ij} the composition $G_n \rightarrow G_n^{\text{ab}} \xrightarrow{\chi_{ij}} \mathbb{Z} \hookrightarrow \mathbb{R}$. We will assume that $n \geq 3$, leaving out G_2 which is isomorphic to a free group of rank 2, which is known to have the R_∞ -property.

We begin by recalling the explicit description of $\Sigma^c(G_n)$ due to Orlandi-Korner [27].

Let $A_{ij} := \mathbb{R}\chi_{ij} + \mathbb{R}\chi_{ji}$, $B_{ijk} := \mathbb{R}(\chi_{ij} - \chi_{kj}) + \mathbb{R}(\chi_{jk} - \chi_{ik}) + \mathbb{R}(\chi_{ki} - \chi_{ji})$, i, j, k pairwise distinct. Note that $A_{ij} = A_{ji}$ and $B_{ijk} = B_{pqr}$ if $\{i, j, k\} = \{p, q, r\}$. Let S be union of vector subspaces $S = \bigcup A_{pq} \cup \bigcup B_{ijk} \subset \text{Hom}(G_n, \mathbb{R})$ where the unions are over all pairs of distinct number $p, q \leq n$ and all pairwise distinct numbers $i, j, k \leq n$. It was shown by Orlandi-Korner [27] that $\Sigma^c(G_n)$ is the image of $S \setminus \{0\} \subset \text{Hom}(G_n, \mathbb{R}) \setminus \{0\}$.

Let \mathcal{S}_n denote the semidirect product $C_2^n \rtimes S_n$ where S_n acts on C_2^n by permuting the coordinates. Here $C_2 = \{1, -1\}$. The group

\mathcal{S}_n acts effectively on F_n the free group with basis $\{x_1, \dots, x_n\}$ where $\pi \in \mathcal{S}_n$ permutes the generators: $\pi(x_j) = x_{\pi(j)}$, $1 \leq j \leq n$ and the action of the k -th factor of C_2^n is given by the automorphism $t_k(x_k) = x_k^{-1}$, $t_k(x_j) = x_j$, $j \neq k$. Thus \mathcal{S}_n is a subgroup of $\text{Aut}(F_n)$. It is readily verified that \mathcal{S}_n normalizes G_n : $t_k \alpha_{i,j} t_k^{-1} = \alpha_{i,j}^{-1}$ if $k = j$ and equals $\alpha_{i,j}$ otherwise; if $\pi \in \mathcal{S}_n$, then $\pi \alpha_{i,j} \pi^{-1} = \alpha_{\pi(i), \pi(j)}$ for all i, j . In particular $\pi^*(A_{ij}) = A_{\pi(i)\pi(j)}$, $\pi^*(B_{ijk}) = B_{\pi(i)\pi(j)\pi(k)}$ for all $\pi \in \mathcal{S}_n$. Thus we have the following

Lemma 3.9. *Let $n \geq 3$. The action of the group $\mathcal{S}_n \subset \text{Aut}(F_n)$ on $\text{Hom}(G_n, \mathbb{R})$ and on $\Sigma^c(G_n)$ is defined by $\pi^*(\chi_{i,j}) = \chi_{\pi(i), \pi(j)}$, $t^*(\chi_{i,j}) = t_i t_j \chi_{i,j}$, for all $\pi \in \mathcal{S}_n$, $t = (t_1, \dots, t_n) \in C_2^n$. \square*

The following proposition is refinement of a statement of Gonçalves and Kochloukova in proof of [15, Theorem 4.11]).

Proposition 3.10. There exists a homomorphism $\eta : \text{Aut}(G_n) \rightarrow \mathcal{S}_n$ which is surjective such that $\phi^*(\chi_{i,j}) = \epsilon_{i,j} \chi_{\sigma(i), \sigma(j)}$, $1 \leq i \neq j \leq n$, where $\epsilon_{i,j} \in \{1, -1\}$ and $\sigma = \eta(\phi) \in \mathcal{S}_n$. In particular, $\text{Aut}(G_n) \cong K \rtimes \mathcal{S}_n$ where $K = \ker(\eta)$.

Proof. Since ϕ^* is a linear isomorphism of $\text{Hom}(G_n, \mathbb{R})$ and since $\phi^* : \Sigma^c(G_n) \rightarrow \Sigma^c(G_n)$ is a homeomorphism, ϕ^* preserves the collections of subspaces $\mathcal{A} := \{A_{ij} \mid 1 \leq i < j \leq n\}$ and $\mathcal{B} := \{B_{ijk} \mid 1 < i < j < k \leq n\}$. Note that \mathcal{B} is non-empty since $n \geq 3$. In our notation A_{pq}, B_{pqr} it is not assumed that $p < q < r$.

It is readily seen that $(A_{pq} + A_{rs}) \cap B_{ijk} = 0$ unless $\{p, q, r, s\} = \{i, j, k\}$. On the other hand $(A_{ij} + A_{ik}) \cap B_{ijk} = \mathbb{R}(\chi_{k,i} - \chi_{j,i})$. It follows that ϕ^* preserves the collection of 1-dimensional spaces $\mathcal{C} := \{\mathbb{R}(\chi_{k,i} - \chi_{j,i}) \mid i, j, k \text{ pairwise distinct}\}$.

Let $\phi^*(A_{ij}) = A_{pq}$, $\phi^*(A_{ik}) = A_{rs}$, where i, j, k are pairwise distinct. Then $\{p, q\} \cap \{r, s\}$ is a singleton, say $s = p$ —so that $\phi^*(A_{ik}) = A_{pr}$ —and $\phi^*(B_{ijk}) = B_{pqr}$. For, otherwise $(A_{ij} + A_{i,k}) \cap B_{ijk}$ is one-dimensional whereas $\phi^*((A_{ij} + A_{ik}) \cap B_{ijk}) = (A_{pq} + A_{pr}) \cap \phi^*(B_{ijk}) = 0$.

In view of the fact that ϕ^* stabilizes \mathcal{C} , we have

$$\phi^*(\chi_{k,i} - \chi_{j,i}) = a(\chi_{r,p} - \chi_{q,p}). \quad (*)$$

On the other hand $\chi_{k,i} \in A_{ik}$ and so $\phi^*(\chi_{k,i}) \in \phi^*(A_{ik}) = A_{pr}$ and so $\phi^*(\chi_{k,i}) = b\chi_{p,r} + c\chi_{r,p}$ for some $b, c \in \mathbb{R}$; similarly $\phi^*(\chi_{j,i}) = b'\chi_{q,p} + c'\chi_{p,q}$ for some $b', c' \in \mathbb{R}$. Therefore

$$\phi^*(\chi_{k,i} - \chi_{j,i}) = b\chi_{p,r} + c\chi_{r,p} - b'\chi_{q,p} - c'\chi_{p,q}. \quad (**)$$

Comparing (*) and (**) we see that $b = 0 = c'$, that is, $\phi^*(\chi_{k,i}) = c\chi_{r,p}$ and $\phi^*(\chi_{j,i}) = b'\chi_{q,p}$. Since $\phi^* : \text{Hom}(G_n; \mathbb{R}) \rightarrow \text{Hom}(G_n, \mathbb{R})$

preserves the lattice $\text{Hom}(G_n, \mathbb{Z})$ and since $\chi_{k,i}, \chi_{j,i}$ are part of a \mathbb{Z} -basis of $\text{Hom}(G_n, \mathbb{Z})$, we see that $c, b' = \pm 1$.

To complete the proof, we define the permutation $\sigma \in S_n$ associated to $\phi \in \text{Aut}(G_n)$ as $\sigma(i) = p$, (with notation as above). Note that σ is indeed a bijection since ϕ^* is an isomorphism. We define $\eta : \text{Aut}(G_n) \rightarrow S_n$ by $\eta(\phi) = \sigma$. Then η is a homomorphism of groups. It is surjective since its restriction to $S_n \subset \mathcal{S}_n$ is the identity by Lemma 3.9. This also shows that η splits, completing the proof. \square

Remark 3.11. It seems plausible that there exists a surjective homomorphism $\tau : \text{Aut}(G_n) \rightarrow \mathcal{S}_n$ such that $\phi^*(\chi_{i,j}) = t_i t_j \chi_{\sigma(i), \sigma(j)}$, $1 \leq i \neq j \leq n$, where $\tau(\phi) = (t_1, \dots, t_n) \in C_2^n$, $\sigma = \eta(\phi) \in S_n$. This would then imply that $\text{Aut}(G_n) \cong N \rtimes \mathcal{S}_n$ for a suitable subgroup $N \subset \text{Aut}(G_n)$.

The above proposition says that the matrix of ϕ^* with respect to the basis $\{\chi_{i,j} \mid 1 \leq i \neq j\}$ (ordered by, say, the lexicographic ordering of the indices i, j), is of the form $\phi^* = DP$ where D is a diagonal matrix with eigenvalues ± 1 and P is a permutation matrix.

Lemma 3.12. *Let $T = DP$ where $D, P \in M_n(\mathbb{R})$ are such that D is a diagonal matrix and P is a permutation matrix. If $P = P_1 \dots P_k$ is a cycle decomposition then there exist eigenvectors v_1, \dots, v_k which are linearly independent.*

Proof. The cycle decomposition allows us to express \mathbb{R}^n as a direct sum $V_1 \oplus \dots \oplus V_k$ where V_j is spanned by $\{e_i \mid P_j(i) \neq i\}$. Specifically, if $P_j = (i_1, \dots, i_k)$. Then $v_j := e_{i_1} + d_{i_1} e_{i_2} + \dots + d_{i_1} \dots d_{i_{k-1}} e_{i_k}$ which is the sum of the vectors in the DP -orbit of e_{i_1} , is an eigenvector of T with eigenvalue $d_{i_1} \dots d_{i_k}$. Evidently v_1, \dots, v_k are linearly independent. \square

We will use the above lemma to construct two linearly independent eigenvectors of ϕ^* (with further properties that will be relevant for our purposes). Let $\sigma = \eta(\phi) \neq id$ and $\phi^* = DP$ with D diagonal and P a permutation transformation (with respect to the basis $\{\chi_{i,j}\}$). Suppose that σ has a k -cycle in its cycle decomposition where $k > 2$. Choose any i that occurs in it and let $j := \sigma(i)$. Then $\chi_{i,j}$ and $\chi_{j,i}$ do not occur in the same orbit of DP and so $v_{i,j} := \sum_{0 \leq r < k} (DP)^r (\chi_{i,j})$ and $v_{j,i} := \sum_{0 \leq j < k} (DP)^r \chi_{j,i}$ are eigenvectors of the same eigenvalue $\epsilon \in \{1, -1\}$. Without loss of generality we assume that $i = 1, j = 2$ and set $v_{1,2} =: u, v_{2,1} = v$. Suppose that there is no such k -cycle in σ . Then σ is a product of disjoint transpositions. Without loss of generality, suppose that the transposition $(1, 3)$ occurs in the decomposition. Since $n > 2$, either σ has a fixed point, say 2, or $n > 3$ and, say the transposition $(2, 4)$ occurs in the decomposition. In the first case

$u := \chi_{1,2} + d_{1,2}\chi_{3,2}$, $v := \chi_{2,1} + d_{2,1}\chi_{2,3}$ are eigenvectors of P and in the latter case, $u := \chi_{1,2} + d_{1,2}\chi_{3,4}$ and $v := \chi_{2,1} + d_{2,1}\chi_{4,3}$ are eigenvectors of P . Thus in *all* case $\chi_{1,2}$ occurs in u and $\chi_{2,1}$ occurs in v where u, v are eigenvectors of ϕ^* . If 1 is an eigenvalue of ϕ^* , then $\bar{\phi}$ has a non-zero fixed element so $R(\phi) = \infty$. So assume that $\phi^*(u) = -u, \phi^*(v) = -v$. Then there exists elements $\beta, \gamma \in G_n$ such that $\bar{\phi}(\beta) = -\beta, \bar{\phi}(\bar{\gamma}) = -\bar{\gamma}$ where $\bar{\alpha}_{1,2}, \bar{\alpha}_{2,1}$ occur in $\bar{\beta}, \bar{\gamma}$ respectively, with coefficient 1.

Denote by $\Gamma_2 := \Gamma_2(G_n)$ the commutator subgroup of G_n and by $\Gamma_3 := \Gamma_3(G_n)$ the subgroup $[G_n, \Gamma_2] \subset \Gamma_2$. Thus G_n/Γ_3 is a two-step-nilpotent group and we have the following exact sequences:

$$\begin{aligned} 1 &\rightarrow \Gamma_3 \rightarrow G_n \rightarrow G_n/\Gamma_3 \rightarrow 1, \\ 1 &\rightarrow \Gamma_2/\Gamma_3 \rightarrow G_n/\Gamma_3 \rightarrow G_n/\Gamma_2 \rightarrow 1 \end{aligned}$$

Since Γ_2 and Γ_3 are characteristic in G_n any automorphism of G_n restricts to automorphisms of Γ_2 and Γ_3 and hence induces automorphisms of the quotients $G/\Gamma_3, \Gamma_2/\Gamma_3$ and $G_n/\Gamma_2 = G_n^{\text{ab}}$.

Denote by $\theta \in \text{Aut}(G_n/\Gamma_3)$ the automorphism defined by ϕ and θ' , the restriction of θ to Γ_2/Γ_3 . With notations as above, $[\beta, \gamma]\Gamma_3 \in \Gamma_2/\Gamma_3$ satisfies $\theta'([\beta, \gamma]\Gamma_3) = [\beta, \gamma]\Gamma_3$. By using the addition formula (Lemma 2.1), we conclude that $R(\theta) = \infty$, *provided* $[\beta, \gamma]\Gamma_3$ is of infinite order. Granting this for the moment, by the first part of the same lemma we conclude that $R(\phi) = \infty$ using the first exact sequence above. Since $\phi \in \text{Aut}(G_n)$ was arbitrary, we conclude that G_n has the R_∞ -property. So all that remains is to show that $[\beta, \gamma]\Gamma_3$ is not a torsion element.

We use the fact that under the surjection $\psi : G_n \rightarrow G_2$ that maps $\alpha_{i,j}$ to $\alpha_{i,j}$ when $\{i, j\} = \{1, 2\}$ and the remaining $\alpha_{i,j}$ to 1, Γ_k maps onto $\Gamma_k(G_2)$, $k = 2, 3$. Let $\beta_2, \gamma_2 \in G_2$ be the images of β, γ respectively under ψ . Then $\bar{\beta}_2 = \bar{\alpha}_{1,2}, \bar{\gamma}_2 = \bar{\alpha}_{2,1} \in G_2^{\text{ab}}$. Therefore $[\beta_2, \gamma_2]\Gamma_3(G_2) = [\alpha_{1,2}, \alpha_{2,1}]\Gamma_3(G_2)$. Since G_2 is a free group with basis $\{\alpha_{1,2}, \alpha_{2,1}\}$ we see that $[\alpha_{1,2}, \alpha_{2,1}]\Gamma_3(G_2)$ generates an infinite cyclic group. Hence the same is true of $[\beta, \gamma]\Gamma_3$. This completes the proof of part (iii) of the main theorem, which is restated below:

Theorem 3.13. *The group $G_n, n \geq 2$, has the R_∞ -property. \square*

4. THE THOMPSON GROUP T

Recall, from §1, the description of the Richard Thompson group T as the group of all orientation preserving piecewise linear homeomorphisms of $\mathbb{S} = I/\{0, 1\}$ with slopes in the multiplicative group generated by $2 \in \mathbb{R}_{>0}$ and break points in $\mathbb{Z}[1/2]$. We regard the Thompson group F as the subgroup of T consisting of elements which fix the element $1 \in \mathbb{S}^1$. In this section we prove the following result.

Theorem 4.1. ([8], [16]). *The Richard Thompson group T has the R_∞ -property.*

The fact that T has the R_∞ property has been proved first by Burillo, Matucci, and Ventura [8] (see also [16]). The crucial point in the proofs of the result above is the same in both [8] and [16] and both the proofs rely on the description of the outer automorphism of T (recalled in Theorem 4.2 below). However, since the approaches before getting to the main point are slightly different, we provide our proof here which may contain some features that are useful for other situations (such as in Remark 4.7 below).

It is readily seen that the reflection map r defined as $r(x) = 1 - x$, $x \in [0, 1]$, induces an automorphism $\rho : T \rightarrow T$ defined as $\rho(f) = r \circ f \circ r^{-1} = r \circ f \circ r$. We now state the following result of Brin.

Theorem 4.2. (Brin [4]) *The group of inner automorphisms of T is of index two in $\text{Aut}(T)$ and the quotient group $\text{Out}(T)$ is generated by ρ .*

As observed in §2.2, for any group Γ and any automorphism $\phi \in \text{Aut}(\Gamma)$, and any $g \in \Gamma$, $R(\phi) = \infty$ if and only if $R(\phi \circ \iota_g) = \infty$. Therefore, to establish the R_∞ -property for Γ , it is enough to show that $R(\phi) = \infty$ for a set of coset representatives of $\text{Out}(\Gamma)$. In the case $\Gamma = T$, in view of Theorem 4.2 due to Brin, we need only show that $R(\rho) = \infty$ and $R(id) = \infty$. The latter equality is established in Proposition 4.5 as an easy consequence of Lemma 4.4 below. Since $\rho^2 = id$, we may apply Remark 2.4 to show that $R(\rho) = \infty$. The main idea is to make use of homeomorphisms in $\text{Fix}(\rho)$, whose supports have arbitrarily large number of disjoint intervals in \mathbb{S}^1 . (This was also the idea used in the proof by Burillo, Matucci, and Ventura [8].)

Definition 4.3. *Let X be a Hausdorff topological space.*

(i) *The support of $f \in \text{Homeo}(X)$ is the open set $\text{supp}(f) := \{x \in X \mid f(x) \neq x\}$.*

(ii) *Let $\sigma : \text{Homeo}(X) \rightarrow \mathbb{N} \cup \{\infty\}$ be defined as follows: $\sigma(id) = 0$, if $f \neq id$, $\sigma(f)$ is the number of connected components of $\text{supp}(f)$ if it is finite, otherwise $\sigma(f) = \infty$.*

Lemma 4.4. *Let $\Gamma \subset \text{Homeo}(X)$ and let σ be as defined above. Suppose that $\theta \in \text{Homeo}(X)$ normalizes Γ . Then $\sigma(f) = \sigma(\theta f \theta^{-1})$.*

Proof. It is clear that the number of connected components of an open set $U \subset X$ remains unchanged under a homeomorphism of X . The lemma follows immediately from the observation that $\text{supp}(\theta f \theta^{-1}) = \theta(\text{supp}(f))$. \square

Proposition 4.5. *The groups F and T have infinitely many conjugacy classes.*

Proof. This follows from Lemma 4.4 on observing that F has elements f with $\sigma(f)$ any arbitrary prescribed positive integer. Since $F \subset T$, the same is true of T as well. \square

Lemma 4.6. *Suppose that $h : \mathbb{R} \rightarrow \mathbb{R}$ is an orientation preserving homeomorphism. Then $\text{supp}(h) = \text{supp}(h^k)$ for any non-zero integer k .*

Proof. Since $\text{supp}(h) = \text{supp}(h^{-1})$ we may assume that $k > 0$. Since h is orientation preserving, it is order preserving. Suppose that $x \in \text{supp}(h)$ so that $h(x) \neq x$. Say, $x < h(x)$. Then applying h to the inequality we obtain $h(x) < h^2(x)$ so that $x < h(x) < h^2(x)$. Repeating this argument yields $x < h(x) < \dots < h^k(x)$ and so $x \in \text{supp}(h^k)$. The case when $x > h(x)$ is analogous. Thus $\text{supp}(h) \subset \text{supp}(h^k)$. On the other hand, if $x \notin \text{supp}(h)$, then $h(x) = x$ and so $h^k(x) = x$ for all k . Therefore equality should hold, completing the proof. \square

We are now ready to prove our main theorem.

Proof of Theorem 4.1: By Theorem 4.2(ii), $\text{Out}(T) \cong \mathbb{Z}/2\mathbb{Z}$ generated by ρ . By Proposition 4.5, $R(\text{id}) = \infty$. It only remains to verify that $R(\rho) = \infty$. We apply Remark 2.4 with $\theta = \rho$, $n = 2$, $\gamma = 1$. It remains to show that $\text{Fix}(\rho)$ has infinitely many elements h such that the h^2 are pairwise non-conjugate.

Let $k \geq 1$. Let $f_k \in F \subset T$ be such that $\text{supp}(f_k) \subset (0, 1/2)$ and has exactly k components. Thus, $\sigma(f_k) = k$. (It is easy to construct such an element.) Then $\text{supp}(\rho(f_k)) = \text{supp}(r f_k r^{-1}) = r(\text{supp}(f_k)) \subset (1/2, 1)$ is disjoint from $\text{supp}(f_k) \subset (0, 1/2)$. In particular $f_k \cdot \rho(f_k) = \rho(f_k) \cdot f_k =: h_k$, $\text{supp}(h_k) = \text{supp}(f_k) \cup r(\text{supp}(f_k))$ and so $\sigma(h_k) = 2k$. Moreover, since $\rho^2 = 1$, we see that $h_k \in \text{Fix}(\rho)$. By Lemma 4.6, we have $\sigma(h_k^2) = \sigma(h_k) = 2k$. It follows that h_k^2 are pairwise non-conjugate in T , completing the proof. \square

Remark 4.7. In the case of the generalized Thompson groups $T_{n,r}$, suppose that $\theta \in \text{Aut}(T_{n,r})$ is a torsion element, say of order m , our method of proof of Theorem 4.1 can be applied to show that $R(\theta) = \infty$. In fact, applying a theorem of McCleary and Rubin [25], to the group $T_{n,r}$, we obtain that the automorphism group of $T_{n,r}$ equals its normalizer in the group of all homeomorphisms of the circle $\mathbb{S}^1 = [0, r]/\{0, r\}$. Let $\theta \in \text{Aut}(T_{n,r})$ and $f \in \mathbb{S}^1$ is such that $\theta(x) = f x f^{-1}$ with $f \in \text{Homeo}(\mathbb{R}/r\mathbb{Z})$. Suppose $f^m = \gamma \in T_{n,r}$ so that θ represents a torsion element of $\text{Out}(T_{n,r})$. If $\gamma = 1$, our method

of proof of Theorem 4.1 can be applied to show that $R(\theta) = \infty$. See [16] for details. However, when $\gamma \neq 1$, it is not clear to us how to find elements of $\text{Fix}(\theta)$ satisfying the hypotheses of Lemma 2.3. Our approach yields no information about automorphisms which represent non-torsion elements in the outer automorphism group. The study of the R_∞ property for the groups $T_{n,r}$ is a work in progress.

5. DIRECT PRODUCT OF GROUPS

It was shown in [15, Theorem 4.8] that if $G = G_1 \times \cdots \times G_n$ where each G_i is a finitely generated group with the property that $\Sigma^c(G_i)$ is a finite set of discrete character classes, not all of them empty, then there exists a finite index subgroup H of $\text{Aut}(G)$ such that $R(\phi) = \infty$ for all $\phi \in H$. Further, when each G_i is a generalized Richard Thompson group $F_{n_i, \infty}$, $n_i \geq 2$, then G itself has the R_∞ -property.

We shall strengthen the above result here. We make use of a result of Meinert (as did Gonçalves and Kochloukova [15]) that describes the Σ -invariant of a direct product which is recalled below. (Meinert's theorem describes the Σ -invariant in the more general setting of a graph product of groups.)

Let $G = G_1 \times \cdots \times G_n$ and let $r_j = rk(G_j^{\text{ab}})$ so that $S(G_j) \cong \mathbb{S}^{r_j-1}$. We will assume that $r_1 \geq 1$. Then $S(G) = \prod_{1 \leq j \leq n} \text{Hom}(G_j, \mathbb{R}) \setminus \{0\} / \sim \cong \mathbb{S}^{r-1}$ and so $S(G) \cong \mathbb{S}^{r-1}$, $r := \sum_{1 \leq j \leq n} r_j$. It is understood that $S(G_j) = \emptyset$ if $r_j = 0$. The sphere $S(G_i)$ is identified with the subspace of $S(G)$ with the set of points with j -th coordinate equal to zero for all $j \neq i$. Observe that $S(G_i) \cap S(G_j) = \emptyset$ if $i \neq j$. In order to emphasise this we shall write $S(S_i) \sqcup S(S_j)$ to denote their union, thought of as subspaces of $S(G)$.

Recall that $\Sigma^c(G)$ denotes the complement of $\Sigma^1(G) \subset S(G)$.

Theorem 5.1. (Meinert [26]) *Let $G = G_1 \times \cdots \times G_n$ be finitely generated and let $r_1 = rk(G_1^{\text{ab}}) > 0$. With the above notations, $\Sigma^c(G)$ equals $\sqcup_{1 \leq j \leq n} \Sigma^c(G_j)$. \square*

We will exploit the fact that any $\phi \in \text{Aut}(G)$ induces a homeomorphism of the character sphere $S(G)$ which preserves its rational structure. Recall that an element $[\chi] \in S(G)$ is called *discrete* (or *rational*) if $\text{Im}(\chi) \subset \mathbb{R}$ is infinite cyclic, equivalently, χ may be chosen to take values in $\mathbb{Q} \subset \mathbb{R}$. The set of rational points in $S(G)$ is denoted $S_{\mathbb{Q}}(G)$. We denote by $D_{\mathbb{Q}}(G)$ the set of isolated rational points in $\Sigma^c(G)$. The set of all limit points of $D_{\mathbb{Q}}(G)$ which are contained in $S_{\mathbb{Q}}(G)$ will be denoted by $L_{\mathbb{Q}}(G)$. Also, we denote by $L(G)$ the set of all limit points of $\Sigma^c(G)$. Since $\Sigma^c(G)$ is closed, $L_{\mathbb{Q}}(G)$ and $L(G)$ are subsets of $\Sigma^c(G)$.

Any homeomorphism of $\Sigma^c(G)$ induced by an automorphism of G maps $D_{\mathbb{Q}}(G), L_{\mathbb{Q}}(G), L(G)$ respectively onto itself.

We are now ready to prove the following theorem. The proof is essentially the same in spirit as that of [15, Theorem 3.3]. See also [15, §4c].

Theorem 5.2. *Suppose that $G = G_1 \times \cdots \times G_n$, $n \geq 1$, is finitely generated and that any one of the following holds: (i) the set $D_{\mathbb{Q}}(G_1)$ is non-empty, finite, and is contained in an open hemisphere and $D_{\mathbb{Q}}(G_j)$ is finite (possibly empty) for $2 \leq j \leq n$, (ii) the set $L_{\mathbb{Q}}(G_1)$ is non-empty, finite, and is contained in an open hemisphere and $L_{\mathbb{Q}}(G_j)$ is a finite set (possibly empty) for $2 \leq j \leq n$, (iii) the set $L(G_1) \cap S_{\mathbb{Q}}(G_1)$ is a non-empty finite set contained in an open hemisphere and the set $L(G_j) \cap S_{\mathbb{Q}}(G_j)$ is finite (possibly empty) for $2 \leq j \leq n$. Then G has the R_{∞} -property.*

Proof. Let $\phi \in \text{Aut}(G)$. We shall show that there exists a discrete character $\lambda \in \text{Hom}(G, \mathbb{R})$ such that $\lambda \circ \phi = \lambda$. By the discussion in §2.3, it follows that $R(\phi) = \infty$ and it follows that G has the R_{∞} -property.

First we suppose that $n = 1$. The theorem, then, is essentially due to Gonçalves and Kochloukova [15]. Let $\phi \in \text{Aut}(G)$ and let $\phi^* : \Sigma^c(G) \rightarrow \Sigma^c(G)$ be the induced map, defined as $\phi^*([\chi]) = [\chi \circ \phi]$. The map $\phi^* : \Sigma^c(G) \rightarrow \Sigma^c(G)$ being a homeomorphism, it maps isolated points to isolated points. Moreover, ϕ^* preserves the set of all rational points in $\Sigma^c(G)$. It follows that $\phi^*(W) = W$ where W is one of the sets $D_{\mathbb{Q}}(G), L_{\mathbb{Q}}(G)$ or $L(G) \cap S_{\mathbb{Q}}(G)$.

In each of the cases (i)-(iii), we see that there is a non-empty finite set of rational character classes $W(G) \subset S_{\mathbb{Q}}(G)$ which is contained in an open hemisphere and is mapped to itself by ϕ^* . Suppose that $[\chi] \in W(G)$ and that the orbit of $[\chi]$ under ϕ^* , namely the set $\{(\phi^*)^j([\chi]) = [\chi \circ \phi^j] \mid j \in \mathbb{N}\}$, has k elements. Then the orbit sum $\lambda := \sum_{0 \leq j < k} \chi \circ \phi^j \in \text{Hom}(G, \mathbb{R})$ is a non-zero discrete character invariant under ϕ^* , as was to be shown.

Now let $n = 2$. By Meinert's theorem (Theorem 5.1) $D_{\mathbb{Q}}(G) = D_{\mathbb{Q}}(G_1) \sqcup D_{\mathbb{Q}}(G_2)$, $L_{\mathbb{Q}}(G) = L_{\mathbb{Q}}(G_1) \sqcup L_{\mathbb{Q}}(G_2)$ and $L(G) = L(G_1) \sqcup L(G_2)$.

Case (i). Let $[\chi] \in D_{\mathbb{Q}}(G_1)$. Consider the ϕ^* -orbit of $[\chi]$, namely, $\{(\phi^k)^*([\chi]) = [\chi \circ \phi^k] \mid k \in \mathbb{Z}\}$. This set is finite since it is contained in $D_{\mathbb{Q}}(G) = D_{\mathbb{Q}}(G_1) \sqcup D_{\mathbb{Q}}(G_2)$, which is finite. Suppose that $[\chi \circ \phi^j], 0 \leq j < q$, are the distinct rational points in the orbit. We claim that the orbit sum $\lambda := \sum_{0 \leq j < q} \chi \circ \phi^j$ is a non-zero character such that $\lambda \circ \phi = \lambda$. To see that $\lambda \in \text{Hom}(G, \mathbb{R})$ is non-zero, note that its

restriction to G_1 is the character $\lambda_J = \sum_{j \in J} \chi \circ \phi^j$ where $J := \{j < q \mid [\chi \circ \phi^j] \in D_{\mathbb{Q}}(G_1)\}$. Since $D_{\mathbb{Q}}(G_1)$ is contained in an open hemisphere, the characters $\chi \circ \phi^j, j \in J$, are in an open half-space of $\text{Hom}(G_1, \mathbb{R})$. Therefore the same is true of their sum, λ_1 , and we conclude that $\lambda \neq 0$. It is clear that $\lambda \circ \phi = \lambda$ since $[\lambda \circ \phi] = [\lambda]$ and λ is rational. As observed in the first para of §2.3, this implies that $R(\phi) = \infty$.

Proof in cases (ii) is almost identical, starting with a $[\chi] \in L_{\mathbb{Q}}(G_1)$. We need only observe that $\phi^*(L_{\mathbb{Q}}(G)) = L_{\mathbb{Q}}(G)$ and that as in case (ii), $L_{\mathbb{Q}}(G) = L_{\mathbb{Q}}(G_1) \sqcup L_{\mathbb{Q}}(G_2)$ is finite. The orbit sum $\lambda := \sum_{0 \leq j < q} \chi \circ \phi^j$ is again a non-zero character which is discrete and satisfies $\lambda \circ \phi = \lambda$. Again we conclude that $R(\phi) = \infty$.

Case (iii). Again we start with a $\chi \in L(G_1) \cap S_{\mathbb{Q}}(G_1)$ and proceed as in case (ii). We leave the details to the reader.

Finally, let $n \geq 3$ be arbitrary. Let $H = G_2 \times \cdots \times G_n$. Again by Meinert's theorem $D_{\mathbb{Q}}(H) = \sqcup_{2 \leq j \leq n} D_{\mathbb{Q}}(G_j)$ and similar expressions hold for $L_{\mathbb{Q}}(H)$ and $L(H) \cap S_{\mathbb{Q}}(H)$. Our hypotheses on G_j implies that one of the sets $D_{\mathbb{Q}}(H), L_{\mathbb{Q}}(H), L(H) \cap S_{\mathbb{Q}}(H)$ is finite depending on case (i), (ii), and (iii) respectively. Since $G = G_1 \times H$, we are now reduced to the situation where $n = 2$, which has just been established. This completes the proof. \square

We conclude the paper with the following examples.

Examples 5.3. (i) Examples of groups with $D_{\mathbb{Q}}(G)$ non-empty, finite, and contained in an open hemisphere are known. These include non-polycyclic nilpotent-by-finite groups of type FP_{∞} , the generalized Richard Thompson groups $F_{n,\infty}$, the double of a knot group K with non-finitely generated commutator subgroup (thus $G \cong K \star_{\mathbb{Z}^2} K$). For details see [15, §4].

(ii) Examples of groups with $D_{\mathbb{Q}}(G)$ and $L_{\mathbb{Q}}(G)$ being finite sets are finite groups, the Houghton groups [6], the pure symmetric automorphism groups [27], finitely generated infinite groups with finite abelianization (which include the generalized Richard Thompson groups $T_{n,r}$; see [6, p. 64]), $\mathbb{Z}^n, n \geq 1$, and the free groups of rank $n \geq 2$. Another class of such groups is provided by [2, Theorem 8.1]. Consider a finitely generated group G which is a subgroup of the group of all orientation preserving PL-homeomorphisms of the interval $[0, 1]$. The group G is said to be *irreducible* if there is no G fixed point in $(0, 1)$. The logarithms of the slopes near the end points 0, 1, define characters $\chi_0, \chi_1 : G \rightarrow \mathbb{R}$ respectively. We recall that two characters λ, χ are independent if $\lambda(\ker(\chi)) = \lambda(G)$ and $\chi(\ker(\lambda)) = \chi(G)$. It was shown in [2, Theorem 8.1] that $\Sigma^c(G) = \{[\chi_0], [\chi_1]\}$ if G is irreducible and

χ_0, χ_1 are independent. (These points may not be in $S_{\mathbb{Q}}(G)$; see [2, p. 470].)

(iii) Let $G = G_1 \times G_2$ where G_1 is a finite product of groups (with G_1 non-trivial) as in example (i) and G_2 , a finite product of groups as in example (ii) above. Then G has the R_{∞} -property. Since there are continuously many pairwise non-isomorphic two generated infinite simple groups, taking G_2 to be any one of them, we obtain a continuous family of groups with R_{∞} -property.

Acknowledgments: We thank Dessislava Kochloukova for pointing out to us the papers [11] and [9] concerning the Houghton groups. We thank the referee for his/her careful reading of our paper and for pointing out to us the paper [20] and references to Theorem 3.2, Corollary 3.3, namely [28] and [12]. The first named author was partially supported by FAPESP-Fundação de Amparo a Pesquisa do Estado de São Paulo, Projeto Temático Topologia Algébrica, Geométrica e Diferencial 2012/24454-8. The second named author was partially supported by Department of Atomic Energy, Government of India, under a XII Plan Project.

REFERENCES

- [1] C. Bleak, A. Fel'shtyn, and D. L. Gonçalves, *Twisted conjugacy classes in R. Thompson's group F*. Pacific Journal of Mathematics **238** no. 1 (2008), 1–6.
- [2] R. Bieri, W. D. Neumann, and R. Strebel, *A geometric invariant of discrete groups*. Invent. Math. **90** (1987), 451–471.
- [3] R. Bieri and R. Strebel, *On groups of PL-homeomorphisms of the real line*. University of Frankfurt preprint, 1985. Second, corrected augmented version, edited by R. Strebel, arxiv:math/1411.2868.
- [4] Matthew G. Brin, *The chameleon groups of Richard J. Thompson: automorphisms and dynamics*. Publ. Math. IHES. **84** (1997), 5–33.
- [5] Matthew G. Brin and Fernando Guzmán, *Automorphisms of generalized Thompson groups*. J. Alg. **203** no. 1 (1998), 285–348.
- [6] Kenneth S. Brown, *Finiteness properties of groups*. Journal of Pure and Applied Algebra **44** (1987), 45–75.
- [7] Kenneth S. Brown, *Trees, valuations, and the Bieri-Neumann-Strebel invariant*. Invent. math. **90** (1987), 479–504
- [8] J. Burillo, F. Matucci, and E. Ventura, *The conjugacy problem for extensions of the Thompson's group F*. arxiv:math/1307.6750v2.
- [9] J. Burillo, S. Cleary, A. Martino, and C. E. Röver, *Commensurations and Metric Properties of Houghton's Groups*. arxiv:math/1403.0026.
- [10] J. W. Cannon, W. J. Floyd, and W. R. Parry, *Introductory notes on Richard Thompson's groups*. Enseign. Math. (2), **42** 3-4 (1996), 215–256.
- [11] Charles Cox, *Twisted conjugacy in Houghton's group*. arxiv:math/1410.7051v1.
- [12] J. D. Dixon and B. Mortimer, *Permutation groups*. Graduate Texts in Mathematics **163**, Springer-Verlag, New York, 1996.

- [13] A. Felshtyn, *The Reidemeister number of any automorphism of a Gromov hyperbolic group is infinite*. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **279** (2001), Geom. i Topol. **6**, 229–240.
- [14] D. L. Gonçalves and K. Dekimpe, *The R_∞ property for free groups, free nilpotent groups and free solvable groups*. Bull. London Math. Soc. (2014). doi: 10.1112/blms/bdu029.
- [15] D. L. Gonçalves and D. Kochloukova, *Sigma theory and twisted conjugacy classes*. Pacific Jour. Math. **247** (2010), 335–352.
- [16] D. L. Gonçalves and P. Sankaran, *Twisted conjugacy in Richard Thompson’s group T* . arxiv:math/1309.2875v2.
- [17] D. L. Gonçalves and P. N. Wong, *Twisted conjugacy classes in exponential growth groups*. Bull. London Math. Soc. **35** (2003), 261–268.
- [18] D. L. Gonçalves and P. N. Wong, *Homogeneous spaces in coincidence theory II*. Forum Math. **17** (2005), 297–313.
- [19] P. R. Heath, *Product formulae for Nielsen numbers of fibre maps*. Pacific Jour. Math. **117** (1985), 267–289.
- [20] P. R. Heath, *Groupoids and relations among Reidemeister and among Nielsen numbers*. Topology and its Applications **181** (2015), 3–33.
- [21] Graham Higman, *Finitely presented infinite simple groups*. Department of Pure Mathematics, Australian National University, Canberra, Notes Pure Math. **8** 1974.
- [22] J. H. Jo, J. B. Lee, and S. R. Lee, *The R_∞ property for Houghton’s groups*. arXiv:math/1412.8767.
- [23] G. Levitt and M. Lustig, *Most automorphisms of a hyperbolic group have very simply dynamics*. Annales Scient. École Normale Supérieure **33** (2000), 507–517.
- [24] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial group theory*. Interscience Publishers, New York, 1966.
- [25] S. McCleary and M. Rubin, *Locally moving groups and the reconstruction problem for chains and circles*. arXiv:math/0510122.
- [26] H. Meinert, *The Bieri-Neumann-Strebel invariant for graph products of groups*. J. Pure Appl. Alg. **103** (1995), 205–210.
- [27] L. A. Orlandi-Korner, *The Bieri-Neumann-Strebel invariant for basis-conjugating automorphisms of free groups*. Proc. Amer. Math. Soc. **128** no. 5 (2000), 1257–1262.
- [28] W. R. Scott, *Group Theory*. Dover, New York, 1987.
- [29] Melanie Stein, *Groups of piecewise linear homeomorphisms*. Trans. Amer. Math. Soc. **332** no. 2 (1992), 477–514.

DEPARTAMENTNO DE MATEMÁTICA - IME, UNIVERSIDADE DE SÃO PAULO,
 CAIXA POSTAL 66.281 - CEP 05314-970, SÃO PAULO - SP, BRASIL
E-mail address: dlgoncal@ime.usp.br

THE INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI,,
 CHENNAI 600113, INDIA
E-mail address: sankaran@imsc.res.in