


Course IMSc, Chennai, India

January-March 2018



The cellular ansatz:
bijective combinatorics and quadratic algebra

Xavier Viennot

CNRS, LaBRI, Bordeaux

www.viennot.org

mirror website

www.imsc.res.in/~viennot

Chapter 4

Trees and tableaux

Ch4c



IMSc, Chennai
April 19, 2018

Xavier Viennot
CNRS, LaBRI, Bordeaux
www.viennot.org



mirror website
www.imsc.res.in/~viennot


From Ch 4b

Jeu de taquin for an increasing word



"jeu de Taquin"











Malvenuto-Reutenauer
product $*$
in $K[S_\infty]$



Loday-Ronco
product $*$





Bijection


increasing
binary
tree

T



permutation

σ



$$\sigma = 7 \ 4 \ 3 \ 8 \ 2 \ 9 \ 5 \ 1 \ 6$$

Bijection

increasing
binary
tree

T

permutation

σ

$$T \xrightarrow{\Pi} \sigma$$

$$\sigma \xrightarrow{\delta} T$$

symmetric order
(or of "vertices
projection")

"déployé"

$$\text{word } w = u m v$$




$$\delta(w) = \delta(u) \text{ } m \text{ } \delta(v)$$


m (unique)
minimum
letter

alternative binary trees

Definition


edge-alternative binary trees






Definition

edge-alternative woods




Definition

alternative




Nadeau (2011)




bijection

edge-alternative
binary tree



edge-alternative binary trees

alternative
binary tree




alternative binary trees

bijection

alternative
binary trees

permutations



$\xrightarrow{\pi}$
symmetric
order
(projection)

$\sigma = (6 \ 1 \ 4 \ 9 \ 3 \ 5 \ 8 \ 2 \ 7)$


$\bar{\delta}(\sigma) =$

$M = \max(\sigma)$

$\sigma = u M v$
 $\bar{\delta}$ alternative "deploy"

"hook length"
formula
same as for
increasing
binary tree

$$\frac{n!}{\prod_x h_x}$$




"hook-length
formula"

$$\frac{n!}{\prod_x h_x}$$


n nb of vertices

product
of size
of sub-trees



nb of increasing binary
tree
for a binary tree **B**

example










"hook-length"
 h_x


$$\frac{9!}{2^2 \cdot 4 \cdot 7 \cdot 9} = 360$$


jeu de taquin
for


edge alternative binary trees and woods


























































alternative binary trees







alternative tableaux

comparison with the
exchange-fusion algorithm





"exchange-fusion" algorithm











7 8 9 1 2 3 4 5 6


7 4 3 8 2 9 5 1 6





bijection alternative tableaux size n


permutations on $(n+1)$


"exchange-fusion" algorithm





































7 4 8 3 2 9 5 1 6




7 4 3 8 2 9 5 1 6

alternative binary trees



"jeu de taquin"


?




"exchange-fusion" algorithm

comparison with the
exchange-fusion algorithm

the twisted symmetric order



"twisted"
symmetric
order



"twisted"
symmetric
order





visit(B)

- if r root is a left son
visit(R), visit(r), visit(L)
- if r root is a right son
visit(L), visit(r), visit(R)


- The root n of the principal binary tree is considered as a "left son"

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 5 & 3 & 2 & 7 & q & 1 & 4 & 6 \end{pmatrix}$$





"twisted"
symmetric
order

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 5 & 3 & 2 & 7 & 9 & 1 & 4 & 6 \end{pmatrix}$$



?




7 4 3 8 2 9 5 1 6



$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 5 & 3 & 2 & 7 & 9 & 1 & 4 & 6 \end{pmatrix}$$

$$\sigma^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 7 & 4 & 3 & 8 & 2 & 9 & 5 & 1 & 6 \end{pmatrix}$$

7 4 3 8 2 9 5 1 6



$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 5 & 3 & 2 & 7 & 9 & 1 & 4 & 6 \end{pmatrix}$$

Def. Genocchi sequence of a permutation

$$\sigma = \sigma(1) \dots \sigma(n)$$


$$G(\sigma) = z_1 \dots z_{n-1}$$

$$z_x = \begin{cases} a & (\text{ascent}) \\ d & (\text{descent}) \end{cases} \quad \begin{matrix} x = \sigma(i) < \sigma(i+1) \\ \text{"value"} \quad \text{"index"} > \sigma(i+1) \end{matrix}$$


$1 \leq x \leq n-1$

convention : $\sigma(n+1) = 0$ ($\sigma(n)$ is a descent)

$$\sigma = \overset{1}{8} \overset{2}{-5} \overset{3}{-3} \overset{4}{-2} \overset{5}{-7} \overset{6}{-9} \overset{7}{-1} \overset{8}{-4} \overset{9}{-6}$$



9







	■		
	■		
■		■	
	■		



$$\sigma^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 7 & 4 & 3 & 8 & 2 & 9 & 5 & 1 & 6 \end{pmatrix}$$



$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 8 & 5 & 3 & 2 & 7 & 9 & 1 & 4 & 6 \end{pmatrix}$$

comparison with the
underlying binary trees















Second version of the bijection



alternative tableaux

alternative binary trees












tree-like tableaux

alternative
tableaux





tree-like
tableaux



Definition Tree-like tableaux
Aval, Bousicault, Nadeau (2011)

Ferret diagram F with cell



empty cell





pointed cell

(i) the bottom left cell is pointed
(called the root cell)

(ii) for every non-root pointed cell c ,
there exist a pointed cell below c
in the same column, or a pointed cell
to its left in the same row,
but not both

(iii) every column and every row
possesses at least one pointed cell











Catalan tree-like tableaux




Two Bijections Catalan alternative tableaux

Binary trees














Non-ambiguous trees



Aval, Boussicault, Bouvel, Silimbani
(2013) (2014)









non-ambiguous trees

in bijection with permutations
with Genocchi sequence


$\underbrace{a \ a \ \dots \ a}_k \ \underbrace{d \ \dots \ d}_l$



tree-like
tableaux





alternative
tableaux




1	2	3	4	
	●			5
	●			6
●		●		7
	●			8
				9


$$\sigma = (\overset{1}{8} - \overset{2}{6} - \overset{3}{5} - \overset{4}{2} - \overset{5}{4} - \overset{6}{9} - \overset{7}{1} - \overset{8}{3} - \overset{9}{7})$$

alternative
tableaux










tree-like
tableaux




alternative
tableaux




$$\sigma = (\overset{1}{8} - \overset{2}{5} - \overset{3}{3} - \overset{4}{2} - \overset{5}{7} - \overset{6}{9} - \overset{7}{1} - \overset{8}{4} - \overset{9}{6})$$


$$\sigma = (\overset{1}{8} - \overset{2}{6} - \overset{3}{5} - \overset{4}{2} - \overset{5}{4} - \overset{6}{9} - \overset{7}{1} - \overset{8}{3} - \overset{9}{7})$$




4	→	3
7	→	4
3	→	5
5	→	6
6	→	7



$$\sigma = (\overset{1}{8} - \overset{2}{5} - \overset{3}{3} - \overset{4}{2} - \overset{5}{7} - \overset{6}{9} - \overset{7}{1} - \overset{8}{4} - \overset{9}{6})$$




interval
(weak) Bruhat order




$$\sigma = (\overset{1}{8} - \overset{2}{6} - \overset{3}{5} - \overset{4}{2} - \overset{5}{4} - \overset{6}{9} - \overset{7}{1} - \overset{8}{3} - \overset{9}{7})$$

hooks formula



The number of non-ambiguous trees having B as underlying binary tree is

$$\frac{k! l!}{\prod_{\text{edge } e} h_e \prod_{\text{edge } e} h_e}$$



non-ambiguous trees

in bijection with permutations
with Genocchi sequence

$\underbrace{a \ a \ \dots \ a}_k \ \underbrace{d \ \dots \ d}_l$

in bijection with permutations
on $(k+l+1)$ such that all exceedances
are exactly $1, 2, \dots, k$.

Ehrenborg, Steingrímsson (2000)

$$\sum_{k=1}^n |s(n, k)| A(k, l) = n^{l-1} n!$$

$$A(k, l) = \sum_{i=1}^k (-1)^{k-i} S(k, i) i! i^{l-1}$$

$s(n, k)$ Stirling 1st permutations, cycles
 $S(k, i)$ Stirling 2nd partitions, block numbers

Ehrenborg, Steingrimsón (2000)

complete non-ambiguous trees

Bessel functions

and heaps of pieces

b_n

number of complete non-ambiguous trees
of size n (number of internal vertices)
satisfies:

$$\sum_{n \geq 1} b_n \frac{t^n}{(n!)^2} = -\log \left(\sum_{n \geq 0} (-1)^n \frac{t^n}{(n!)^2} \right)$$

A002190

1, 1, 4, 33, 456, 9460

Carlitz (1963)
a sequence of integers related to the
Bessel functions

bijection


complete non-ambiguous trees size $(2n-1)$



pyramids Q_n having n elements

of the form $(i, \sigma(i))$ where $\sigma \in S_n$
and the unique maximal element
is $(1, \sigma(1))$


Emma Jin (2016)


heaps of "sticks"






heaps of "sticks"












$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 3 & 4 & 1 & 2 \end{pmatrix}$$

heaps of "sticks"





Aisbett (2015)

poset of vector partitions of $[n]$
into s components $\Pi_{n,s}$

CL-shellable

$\Rightarrow$ homotopy type of
wedge of spheres
dimension $(n-2)$


$s=1$, number of
spheres $= b_n$

Carlitz, Scoville, Vaughan (1974)

number of pairs of permutations
of $\{1, \dots, n\}$ with no common rise

Beáta Bényi, G. Nagy
(2017)

Fedou, Rawlings (1995)



q -Polya enumeration

parallelogram
polyominoes

($area$, number of rows
of columns)

= staircase polygons

Delest, Fedou (1990)


$$\frac{J_1(t; q)}{J_0(t; q)}$$

q -Bessel


heaps of "segments"

Bousquet-Mélou, X.V. (1992)

staircase
polygon



$a_{m,n,p}$



poids

$q^i t u^{(j-i)}$

heaps of "segments"

Bousquet-Mélou, X.V. (1992)

heaps of "sticks"

Fedou (1992)

Bijection


parallelogram polyominoes
(=staircase polygons)


Binary trees


via non-ambiguous trees


parallelogram
polyominoes


= staircase polygons













parallelogram
polyominoes

bijection

binary trees



Aval, Boussicault, Bouvel, Silimbani
(2013) (2014)

Aval, Boussicault, Delcroix-Oger,
Hivert, Laborde-Zubieta (2015)

Reverse Q-tableaux

example with the PASEP algebra:

tree-like tableaux

PASEP algebra

$$Q \left\{ \begin{array}{l} DE = q ED + EX + YD \\ XE = EX \\ DY = YD \\ XY = YX \end{array} \right.$$

$$Q \left\{ \begin{array}{l} DE = \square ED - \blacksquare EX - \blacksquare YD \\ XE = \square EX \\ DY = \square YD \\ XY = \square YX \end{array} \right.$$

$$\frac{D}{X} \quad Y|E$$

alternative
tableaux

PASEP algebra

Q $\left\{ \begin{array}{l} DE = q ED + EX + YD \\ XE = EX \\ DY = YD \\ XY = YX \end{array} \right.$

The diagram illustrates the derivation of the first equation in the set. The terms EX and YD in the first equation are circled in blue. Arrows point from these circles to the corresponding terms in the equations $XE = EX$ and $DY = YD$, indicating that these terms are defined or simplified by those equations.

PASEP algebra

Q {

$$\begin{aligned} DE &= \square ED + \blacksquare EX + \blacksquare YD \\ XE &= \square EX \\ DY &= \square YD \\ XY &= \square YX \end{aligned}$$

The diagram illustrates the derivation of the first equation in the PASEP algebra. The first equation is $DE = \square ED + \blacksquare EX + \blacksquare YD$. The terms EX and YD are circled in blue. Arrows point from these circles to the corresponding terms in the second and third equations: $XE = \square EX$ and $DY = \square YD$. The term ED in the first equation has a small orange '1' written below it.

$$Q \left\{ \begin{array}{l} DE = \square ED - \blacksquare EX - \blacksquare YD \\ XE = \square EX \\ DY = \square YD \\ XY = \square YX \end{array} \right.$$

PASEP algebra

$$\frac{D}{X} \quad Y|E$$

alternative
tableaux


reverse PASEP algebra

$$Q^+ \left\{ \begin{array}{l} DE = \square ED - \blacksquare EX + \blacksquare YD \\ XE = \square EX \\ DY = \square YD \\ XY = \square YX \end{array} \right.$$


$$\frac{D}{X} \quad Y|E$$

tree-like
tableaux


alternative
tableaux




reverse Q-tableaux




reverse Q-tableaux




reverse Q-tableaux






tree-like
tableaux

alternative
tableaux




tree-like
tableaux




tree-like tableaux
are "the" reverse Q -tableaux
of alternative tableaux


tree-like
tableaux



tree-like
tableaux



tree-like
tableaux



In conclusion

The philosophy of the cellular ansatz

The philosophy of the cellular ansatz

combinatorial representation
of the quadratic algebra

$$UD = qDU + Id$$



RSK

~ commutations diagrams

Fomin growth diagrams




~ local rules on edges

RSK

~ duplication of equations



RSK

$$UD = DU + I_v I_h$$










$$U\mathcal{D} = \mathcal{D}U + I_v I_h$$








"commutation diagrams"




bijection





Fomin growth diagrams





"local rules"
on the **edges**



$$\begin{cases} U \mathcal{D} = \mathcal{D} U + YX \\ U Y = Y U \\ X U = U X \\ X Y = YX \end{cases}$$

$$\mathcal{D} \begin{array}{|c|} \hline U \\ \hline \diagup \\ \hline \end{array} \mathcal{D} \quad Y \begin{array}{|c|} \hline U \\ \hline \diagup \\ \hline X \\ \hline \end{array} \mathcal{D}$$

duplication of the commutation relations defining the quadratic algebra Q

$$U \mathcal{D} = \mathcal{D} U + \gamma_1 X_1$$

$$X_1 \gamma_1 = \gamma_2 X_2$$

$$X_2 \gamma_2 = \gamma_3 X_3$$

$$\dots$$


$$X_i \gamma_i = \gamma_{i+1} X_{i+1}$$

$$\dots$$



$$U \gamma_i = \gamma_i U$$

$$X_j U = U X_j$$

$$\dots$$



"local rules"
on the edges





Weil-Heisenberg algebra

$$Q \begin{cases} UD = q_1 DU + t \boxed{YX} \\ UY = YU \\ XD = DX \\ XY = q_2 \boxed{YX} \end{cases}$$



for the Weyl-Heisenberg algebra
 $Q^+ \supset Q$

$$Q \left\{ \begin{array}{l} UD = q_1 DU + t YX \\ UY = YU \\ XD = DX \\ XY = q_2 YX \end{array} \right.$$



$$Q^+ \left\{ \begin{array}{l} YX = q_2 XY + t XY \\ YU = UY \\ DX = XD \\ DU = q_1 UD \end{array} \right.$$



duplication of equations
in the reverse quadratic algebra Q^+



"local rules"
on the edges



The philosophy of the cellular ansatz

combinatorial representation
of the quadratic algebra

$$UD = qDU + Id$$



RSK

~ commutations diagrams

Fomin growth diagrams

~ local rules on edges

RSK

~ duplication of equations


RSK

The philosophy of the cellular ansatz

combinatorial representation
of the quadratic algebra

$$\mathcal{D}E = qED + E + \mathcal{D}$$

alternative
tableaux






- commutations diagrams

$$EXF$$


"Laguerre histories"

- duplication of equations




Adela bijection




"commutation diagrams"



$$AS = SA + J + K$$




$$AS = SA + I_v J + K I_h$$



4 1 6 9 7 8 3 5 2

Laguerre histories



Prop.

T

alternative
tableau



"exchange - fusion"
inverse algorithm

τ

"local"
algorithm

from $D E = E D + E + D$

$$\sigma = \tau^{-1}$$

PASEP algebra

Q


$$\begin{cases} DE = qED + EX + YD \\ XE = EX \\ DY = YD \\ XY = YX \end{cases}$$

$$DE = ED + EX_1 + Y_1 D$$

$$\begin{cases} X_1 E = E X_2 \\ \dots \\ X_i E = E X_{i+1} \\ \dots \end{cases}$$


$$\begin{cases} D Y_1 = Y_2 D \\ \dots \\ D Y_i = Y_{i+1} D \\ \dots \end{cases}$$

$$X_i Y_j = Y_j X_i$$



Adela bijection

Adela $(T) = (P, Q)$




The philosophy of the cellular ansatz

combinatorial representation
of the quadratic algebra

$$\mathcal{D}E = qED + E + \mathcal{D}$$

alternative
tableaux



- commutations diagrams

$$EXF$$


"Laguerre histories"

- duplication of equations

Adela bijection

- local rules on edges ?

with duplication of equations
in the reverse quadratic algebra



PASEP algebra

$$Q \left\{ \begin{array}{l} DE = \boxed{} ED - \boxed{} EX - \boxed{} YD \\ EX = \boxed{} EX \\ DY = \boxed{} YD \\ XY = \boxed{} YX \end{array} \right. \quad \frac{D}{X} \quad Y|E$$

reverse PASEP algebra

$$Q^+ \left\{ \begin{array}{l} ED = \boxed{} \textcircled{DE} \\ EX = \boxed{} XE + \boxed{} \textcircled{DE} \\ YD = \boxed{} DY \\ YX = \boxed{} XY \end{array} \right. \quad \begin{array}{l} \boxed{} \textcircled{DE} \\ \boxed{} \textcircled{DE} \end{array}$$

$$AS = SA + I_v J + K I_h$$

$$AK = KA + I_v A$$

$$JS = SJ + S I_h$$

$$JK = KJ$$

$$A I_v = I_v A$$

$$J I_v = I_v J$$

$$I_h S = S I_h$$

$$I_h K = K I_h$$

$$D = A + J$$

$$E = S + K$$

$$I_v = Y$$

$$I_h = X$$

duplication of equations
in the reverse quadratic algebra Q^+

$$D \rightarrow \begin{cases} A \\ J \end{cases} \quad E \rightarrow \begin{cases} S \\ K \end{cases}$$

$$\begin{array}{c} y \mid \\ \hline A \\ S \mid \\ \hline X \end{array} = \begin{array}{c} A \\ \mid \\ \text{blue square} \\ \mid \\ J \\ \mid \\ \text{red square} \\ \mid \\ \downarrow \end{array} \mid \begin{array}{c} K \\ S \end{array}$$

$$\begin{array}{c} S \\ K \end{array} \mid \begin{array}{c} \text{orange square} \\ \mid \\ A, J \end{array} = \begin{array}{c} A, J \\ \mid \\ S \\ K \end{array}$$

duplication of equations
in the reverse quadratic algebra Q^+

$$\begin{array}{c}
 Y \mid \underline{\quad} \\
 \quad A \\
 S \mid \underline{\quad} \\
 \quad X
 \end{array}
 =
 \begin{array}{c}
 \overline{A} \\
 \leftarrow \text{blue square} \mid K \\
 \overline{J} \\
 \text{red square} \mid S \\
 \downarrow
 \end{array}$$

$$\begin{array}{c}
 Y \mid \underline{\text{blue square}} \\
 \quad A_i \\
 S_i \mid \underline{\text{red square}} \\
 \quad \downarrow \\
 \quad X
 \end{array}
 =
 \begin{array}{c}
 \overline{A_i} \\
 \mid K_i \\
 \overline{J_i} \\
 \mid S_i
 \end{array}
 \qquad
 \begin{array}{c}
 Y \mid \underline{\text{blue square}} \\
 \quad J_i \\
 K_i \mid \underline{\text{red square}} \\
 \quad \downarrow \\
 \quad X
 \end{array}
 =
 \begin{array}{c}
 \overline{A_i} \\
 \mid S_i \\
 \overline{A_i} \\
 \mid S_{i+1}
 \end{array}$$

$$\begin{array}{c} S \\ K \end{array} \bigg| \begin{array}{c} \square \\ \hline A, J \end{array} = \begin{array}{c} A, J \\ \hline S \\ K \end{array}$$

$\begin{array}{c} S_j \\ \hline A_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} A_i \\ \hline \end{array} \bigg \begin{array}{c} S_{j+1} \\ \hline \end{array} \quad (i < j)$	$\begin{array}{c} S_j \\ \hline A_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} A_{i+1} \\ \hline \end{array} \bigg \begin{array}{c} S_j \\ \hline \end{array} \quad (j \leq i)$
$\begin{array}{c} K_j \\ \hline A_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} A_i \\ \hline \end{array} \bigg \begin{array}{c} K_{j+1} \\ \hline \end{array} \quad (i \leq j)$	$\begin{array}{c} K_j \\ \hline A_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} A_i \\ \hline \end{array} \bigg \begin{array}{c} K_j \\ \hline \end{array} \quad (j < i)$

$\begin{array}{c} S_j \\ \hline J_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} J_i \\ \hline \end{array} \bigg \begin{array}{c} S_j \\ \hline \end{array} \quad (i < j)$	$\begin{array}{c} S_j \\ \hline J_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} J_{i+1} \\ \hline \end{array} \bigg \begin{array}{c} S_j \\ \hline \end{array} \quad (j \leq i)$
$\begin{array}{c} K_j \\ \hline J_i \end{array} \bigg \begin{array}{c} \square \\ \hline \end{array} = \begin{array}{c} J_i \\ \hline \end{array} \bigg \begin{array}{c} K_j \\ \hline \end{array}$	

$$\begin{array}{c}
 y \mid \leftarrow \frac{A_i, J_j}{A_i, J_j} = \frac{A_i, J_j}{\mid} y \\
 \hline
 K_j, S_i \mid \frac{x}{x} = \frac{x}{\mid} K_j, S_i
 \end{array}$$


$$y \mid \frac{x}{x} = \frac{x}{\mid} y$$

The philosophy of the cellular ansatz

combinatorial representation
of the quadratic algebra

$$\mathcal{D}E = qED + E + \mathcal{D}$$

alternative
tableaux



~ commutations diagrams

EXF


"Laguerre histories"

Equivalence




~ local rules on edges


with duplication of equations
in the reverse quadratic algebra




K_0




Y



K_0



K_0





J_0

X

J_0

X





"commutation diagrams"

Laguerre histories

A new bijection

demultiplication of equations
in the
reverse PASEP quadratic algebra

(2)

PASEP algebra

$$Q \left\{ \begin{array}{l} DE = \square ED - \square EX + \square YD \\ EX = \square EX \\ DY = \square YD \\ XY = \square YX \end{array} \right.$$

$$\frac{D}{X} \quad Y|E$$

reverse PASEP algebra


$$Q^+ \left\{ \begin{array}{l} ED = \square DE \\ EX = \square XE + \square DE \\ YD = \square DY + \square DE \\ YX = \square XY \end{array} \right.$$

reverse PASEP algebra

$$Q^+ \begin{cases} ED = q \circ DE \\ EX = XE + \circ DE \\ YD = DY + \circ DE \\ YX = XY \end{cases}$$

$$\begin{cases} E_i D_j = q \circ D_{j+1} E_{i+1} \\ E_i X = X E_i + \circ D_1 E_{i+1} \\ Y D_j = D_j Y + \circ D_{j+1} E_1 \\ YX = XY \end{cases}$$

$$i, j \geq 1$$





The Tamil bijection


alternative tableaux (size n)
or tree-like

some words $w \in \{D_i, E_j; i, j \geq 1\}^*$
 $|w| = n$


$$w = D_1 D_5 E_1 D_2 E_2 E_4 D_2 E_2$$









The Tamil bijection




$$w = D_1 D_5 E_1 D_2 E_2 E_4 D_2 E_2$$


Catalan
tableaux










$$w = D_2 D_1 E_2 E_2 D_2 D_1 E_2 E_2 D_1$$




example











The Tamil bijection

binary trees




The Tamil bijection

binary trees



$w = 2\ 1\ 2\ 2\ 1\ 3\ 1\ 1\ 2\ 2\ 1\ 4\ 1\ 1\ 2\ 1$



The ultimate end of the bijective course III

Thank you very much !

for all of you, students, professors, friends,

For the videos:
Gayathri and Kirubananth

special thanks to Amri Prasad





ॐ सरस्वत्यै नमः।